

# **ACTIMMUNE**

---

## **MEDICATION(S)**

ACTIMMUNE

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

1. Hypersensitivity to Actimmune or E. coli derived products

## **REQUIRED MEDICAL INFORMATION**

1. Dx chronic granulomatous disease OR Dx severe malignant osteopetrosis

## **AGE RESTRICTION**

N/A

## **PRESCRIBER RESTRICTION**

N/A

## **COVERAGE DURATION**

Plan Year

## **OTHER CRITERIA**

BvD

## **AKEEGA**

---

### **MEDICATION(S)**

AKEEGA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of deleterious or suspected deleterious BRCA-mutated (BRCAm) metastatic castration-resistant prostate cancer (mCRPC) 2. Contraindication, intolerance, or failure of Lynparza

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ALECENSA**

---

### **MEDICATION(S)**

ALECENSA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of NSCLC that is ALK-positive    a. Tumors are anaplastic lymphoma kinase (ALK)-positive

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

# ALS

---

## **MEDICATION(S)**

RADICAVA ORS, RADICAVA ORS STARTER KIT

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

N/A

## **REQUIRED MEDICAL INFORMATION**

1)Dx of amyotrophic lateral sclerosis (ALS) as defined by the revised El Escorial criteria

## **AGE RESTRICTION**

N/A

## **PRESCRIBER RESTRICTION**

N/A

## **COVERAGE DURATION**

Plan Year

## **OTHER CRITERIA**

N/A

## **ALUNBRIG**

---

### **MEDICATION(S)**

ALUNBRIG

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1)Dx of metastatic non-small cell lung cancer (NSCLC) a)Tumors are anaplastic lymphoma kinase (ALK)-positive 2)Patient has failed/intolerance/contraindication to Alecensa

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ANTIPSYCHOTIC**

---

### **MEDICATION(S)**

CAPLYTA, COBENFY, COBENFY STARTER PACK, FANAPT, FANAPT TITRATION PACK A, LYBALVI, OPIPZA, VERSACLOZ, VRAYLAR 1.5 MG CAP, VRAYLAR 3 MG CAP, VRAYLAR 4.5 MG CAP, VRAYLAR 6 MG CAP

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Previous trial on at least ONE of the following: aripiprazole, clozapine, fluoxetine-olanzapine, haloperidol, olanzapine, quetiapine, risperidone, ziprasidone 2. For Major Depressive Disorder (MDD) or Schizophrenia: Previous trial on Rexulti

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **APO B**

---

### **MEDICATION(S)**

JUXTAPID

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Pt has none of the following health conditions or health concerns: a. History of significant hepatic disease, b. Alcohol abuse,

### **REQUIRED MEDICAL INFORMATION**

1. Pt has untreated, fasting LDL cholesterol greater than 500 mg/dL AND triglycerides less than 300 mg/dL, 2. Pt meets a OR b AND c of the following: a. Pt has documented mutations in both alleles of the LDL receptor or of other genes known to affect LDL receptor function, b. Both of pt's parents have a hx of untreated total cholesterol of greater than 250 mg/dL, c. Pt has xanthomas present before age 10, 3. Pt has failed or is currently taking at least ONE of the following: a. Atorvastatin, Rosuvastatin, or Simvastatin b. Has documented intolerance (e.g. rhabdomyolysis) to statin therapy 4. Pt has previous trial of Repatha OR Praluent

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

1. Initial: 6 mo, 2. Reauthorization: Plan Year

### **OTHER CRITERIA**

N/A

## **AQNEURSA**

---

### **MEDICATION(S)**

AQNEURSA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Use in combination with Miplyffa

### **REQUIRED MEDICAL INFORMATION**

1. Pt has tried and failed at least 3 months of therapy with miglustat as evidenced by a lack of improvement in disease progression (e.g. horizontal saccadic eye movements, SARA scale, R4DNPCCSS score) 2. Pt weighs greater than or equal to 15kg 3. Dx is genetically confirmed (deoxyribonucleic acid [DNA] sequence analysis) by mutations in both alleles of NPC1 or NPC2 OR if there is a mutation in only one allele of NPC1 or NPC2, pt has positive filipin staining or elevated cholestane triol/oxysterols (greater than 2x upper limit of normal) 4. Pt is presenting with at least one neurological symptom of the disease (for example, but not limited to, hearing loss, vertical supranuclear gaze palsy, ataxia, dementia, dystonia, seizures, dysarthria, or dysphagia) 5. Pt is able to walk either independently or with assistance 6. Pt has a SARA score of greater than or equal to 7 and less than or equal to 34 points (out of 40) AND one of the following: A. Within in 2-7 range (0-8 range) of the Gait subtest of the SARA scale B. Be able to perform the 9-Hole Peg Test with Dominant Hand ((HPT-D) (SCAFI subtest) in greater than or equal to 20 to less than or equal to 150 seconds 7. Reauthorization A. fSARA score has remained stable/improved

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Neurologist

### **COVERAGE DURATION**

6 months



**OTHER CRITERIA**

N/A

## **ARCALYST**

---

### **MEDICATION(S)**

ARCALYST

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Combination use with a TNF-inhibitor

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of cryopyrin-associated periodic syndrome 2. Diagnosis of recurrent pericarditis 3. Diagnosis of deficiency of Interleukin-1 Receptor Antagonist (DIRA)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

BvsD Determination

## **ATOPIC DERM**

---

### **MEDICATION(S)**

ADBRY, CIBINQO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Used in combination with another biologic (e.g., Dupixent, Xolair, Humira, Entyvio, and Otezla, etc.)

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of moderate to severe atopic dermatitis (AD) A. Greater than or equal to 10 percent body surface area coverage B. Failure of two of the following: i. Topical corticosteroid (e.g., fluocinonide, clobetasol, etc.) ii. Topical calcineurin inhibitor (eg. tacrolimus ointment 0.1%) iii. Phototherapy iv. Oral immunomodulator (azathioprine, cyclosporine, or mycophenolate) v. Topical PDE-4 (Eucrisa) 2. For reauthorization of Adbry A. a) For patients less than 100kg who achieved clear or almost clear skin, every 4 week dosing has been tried

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## AUGTYRO

---

### MEDICATION(S)

AUGTYRO

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Dx of locally advanced or metastatic ROS1-positive non-small cell lung cancer (NSCLC) or NTRK gene fusion-positive solid tumors 2. Contraindication, intolerance, or failure of Rozlytrek

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A

## **AUSTEDO**

---

### **MEDICATION(S)**

AUSTEDO, AUSTEDO XR, AUSTEDO XR PATIENT TITRATION

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of chorea associated with Huntington's disease 2. Dx of tardive dyskinesia

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **AVMAPKI FAKZYNJA**

---

### **MEDICATION(S)**

AVMAPKI FAKZYNJA CO-PACK

**PENDING CMS APPROVAL**

## **AYVAKIT**

---

### **MEDICATION(S)**

AYVAKIT

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1)Diagnosis of: a)Unresectable or metastatic GIST that is platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation positive including PDGFRA D842V mutations. b)Advanced Systemic Mastocytosis: AdvSM includes patients with aggressive systemic mastocytosis (ASM), systemic mastocytosis with an associated hematological neoplasm (SM-AHN), and mast cell leukemia (MCL). c)Indolent Systemic Mastocytosis

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **BALVERSA**

---

### **MEDICATION(S)**

BALVERSA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of locally advanced or metastatic urothelial carcinoma 2. Confirmed fibroblast growth factor receptor (FGFR3) genetic alteration 3. Progression on or after at least one line of prior systemic therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **BANZEL**

---

### **MEDICATION(S)**

RUFINAMIDE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has inadequate seizure control despite treatment with at least ONE anti-epileptic drug

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **BENLYSTA**

---

### **MEDICATION(S)**

BENLYSTA 200 MG/ML SOLN A-INJ, BENLYSTA 200 MG/ML SOLN PRSYR

### **PENDING CMS APPROVAL**

## **BESREMI**

---

### **MEDICATION(S)**

BESREMI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Polycythemia Vera 2. Inadequate response or intolerance to hydroxyurea

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: 6 months Reauth: Plan Year

### **OTHER CRITERIA**

N/A

## **BOSULIF**

---

### **MEDICATION(S)**

BOSULIF

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt is diagnosed with Philadelphia chromosome positive (Ph+) CML a. Pt's CML is newly diagnosed in chronic phase b. Pt's CML is in chronic phase, accelerated phase, or blast phase, 2. For CML in chronic phase, accelerated phase, or blast phase: a. Pt has previous failure or intolerance to imatinib

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **BRAFTOVI**

---

### **MEDICATION(S)**

BRAFTOVI 75 MG CAP, MEKTOVI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of BRAF V600E or V600K mutation-positive unresectable or metastatic melanoma a. Will be used as combination therapy with encorafenib [Braftovi] and binimetinib [Mektovi] 2. For encorafenib [Braftovi], dx of BRAF V600E mutation-positive metastatic colorectal cancer a. Will be used as combination therapy with cetuximab [Erbix] 3. Dx of BRAF V600E mutation-positive metastatic non-small cell lung cancer (NSCLC) a. Will be used as combination therapy with encorafenib [Braftovi] and binimetinib [Mektovi]

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **BRONCHITOL**

---

### **MEDICATION(S)**

BRONCHITOL, BRONCHITOL TOLERANCE TEST

### **PENDING CMS APPROVAL**

# **BRUKINSA**

---

## **MEDICATION(S)**

BRUKINSA 80 MG CAP

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

N/A

## **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of
  - A. Mantle cell lymphoma (MCL)
  - B. Marginal zone lymphoma (MZL)
  - C. Waldenstrom macroglobulinemia (WM)
  - D. Chronic lymphocytic leukemia (CLL)/ Small lymphocytic lymphoma (SLL)
  - E. Relapsed or refractory follicular lymphoma (FL)
2. For diagnosis of MCL:
  - A. Patient has tried and failed or has contraindication to both Calquence AND Imbruvica
3. For diagnosis of MZL:
  - A. Patient has been previously treated with anti-CD20-based regimen
4. For diagnosis of WM:
  - A. Patient has tried and failed or has contraindication to Imbruvica
5. For diagnosis of CLL/SLL:
  - A. Patient has tried and failed or has contraindication to both Calquence AND Imbruvica
6. For diagnosis of FL:
  - A. Patient has received two or more lines of therapy and Brukinsa will be used in combination with obinutuzumab

## **AGE RESTRICTION**

N/A

## **PRESCRIBER RESTRICTION**

N/A

**COVERAGE DURATION**

Plan Year

**OTHER CRITERIA**

N/A



## **CABLIVI**

---

### **MEDICATION(S)**

CABLIVI

**PENDING CMS APPROVAL**

## **CABOMETYX**

---

### **MEDICATION(S)**

CABOMETYX

**PENDING CMS APPROVAL**

## **CALQUENCE**

---

### **MEDICATION(S)**

CALQUENCE 100 MG TAB

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of mantle cell lymphoma a. Pt has been treated with at least one prior therapy 2. Dx of chronic lymphocytic leukemia 3. Dx of small lymphocytic lymphoma

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **CAPRELSA**

---

### **MEDICATION(S)**

CAPRELSA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx symptomatic or progressive medullary thyroid cancer with unresectable locally advanced or metastatic disease

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **CARBAGLU**

---

### **MEDICATION(S)**

CARGLUMIC ACID, SAPROPTERIN DIHYDROCHLORIDE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For carglumic acid: hyperammonemia due to N-acetylglutamate synthase deficiency 2. For sapropterin: hyperphenylalaninemia due to tetra hydrobiopterin- (BH4-) responsive Phenylketonuria

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **CHOLBAM**

---

### **MEDICATION(S)**

CHOLBAM

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Initial: a. Pt has abnormal results from a urinary bile acids analysis by FAB-MS and neurologic exam,
2. Reauth: a. Patient has experienced improvement in ALT/AST values, bilirubin values, and/or weight

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: Plan year Reauth: Plan Year

### **OTHER CRITERIA**

N/A

## **COMETRIQ**

---

### **MEDICATION(S)**

COMETRIQ (100 MG DAILY DOSE), COMETRIQ (140 MG DAILY DOSE), COMETRIQ (60 MG DAILY DOSE)

**PENDING CMS APPROVAL**

## **COPIKTRA**

---

### **MEDICATION(S)**

COPIKTRA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

Previous use of 2 prior therapies for indication

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **CRESEMBA**

---

### **MEDICATION(S)**

CRESEMBA 186 MG CAP, CRESEMBA 74.5 MG CAP

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Treatment initiated incident to a hospitalization
2. Pt has confirmed fungal infection with one of the following:
  - a. Invasive aspergillosis
  - b. Invasive mucormycosis

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Infectious disease specialist

### **COVERAGE DURATION**

6 months

### **OTHER CRITERIA**

N/A

## **CRINONE**

---

### **MEDICATION(S)**

CRINONE 4 % GEL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

A. Diagnosis of secondary amenorrhea

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **CYSTAGON**

---

### **MEDICATION(S)**

CYSTAGON

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of nephropathic cystinosis 2. Elevated baseline WBC cysteine levels greater than 2 nmol per 1/2 cystine/mg protein 3. CTNS gene mutation 4. Clinical symptoms of an electrolyte imbalance and polyuria

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **CYSTIC FIBROSIS**

---

### **MEDICATION(S)**

CAYSTON, TOBRAMYCIN 300 MG/5ML NEBU SOLN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has had at least ONE positive culture for *Pseudomonas aeruginosa* 2. If request not for generic tobramycin: Previous trial on generic tobramycin

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

BvsD determination for tobramycin only, no BvD for Cayston

## **DANZITEN**

---

### **MEDICATION(S)**

DANZITEN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of: a. newly diagnosed Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase b. chronic phase (CP) and accelerated phase (AP) Ph+ CML i. Documented resistance, intolerance, or contraindication to imatinib (Gleevec)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## DAURISMO

---

### MEDICATION(S)

DAURISMO

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Equal to or greater than 75 years or has comorbidity preventing use of intensive induction chemotherapy. 2. Be given in combination with low-dose cytarabine

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A

## **DIACOMIT**

---

### **MEDICATION(S)**

DIACOMIT

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Dravet syndrome 2. Previous use of clobazam and valproic acid 3. To be used in combination with clobazam

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **DOPTELET**

---

### **MEDICATION(S)**

DOPTELET

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Platelet count less than 50,000 2. Pt is scheduled for a procedure where there is a bleeding risk 3. Doptelet will be used for 5 days starting 10 to 13 days prior to the procedure and discontinued 5 to 8 days prior to the procedure 4. For chronic immune thrombocytopenia, has the patient had an insufficient response to a previous treatment (e.g. corticosteroid, immune globulin)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **DRONABINOL**

---

### **MEDICATION(S)**

DRONABINOL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For HIV-associated wasting syndrome OR cancer-associated anorexia dx: a. Pt has previous trial on megestrol, 2. For CINV dx: a. Pt has previous trial on olanzapine

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

BvD determination

## **DUPIXENT**

---

### **MEDICATION(S)**

DUPIXENT 300 MG/2ML SOLN A-INJ, DUPIXENT 300 MG/2ML SOLN PRSYR

### **PENDING CMS APPROVAL**

## **EMGALITY**

---

### **MEDICATION(S)**

EMGALITY, EMGALITY (300 MG DOSE)

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1.Diagnosis of episodic cluster headache 2. Diagnosis of Episodic Migraine (4-14 migraine days per month) or Chronic Migraine (greater than 14 migraine days per month) A. Member has tried Ajovy  
Reauthorization Criteria: 1. Patient has had a reduction in the number of migraine days per month

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ENBREL**

---

### **MEDICATION(S)**

ENBREL 25 MG/0.5ML SOLN PRSYR, ENBREL 25 MG/0.5ML SOLUTION, ENBREL 50 MG/ML SOLN PRSYR, ENBREL MINI, ENBREL SURECLICK

### **PENDING CMS APPROVAL**

## ENDARI

---

### MEDICATION(S)

L-GLUTAMINE 5 GM PACKET

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

1. Renal insufficiency, 2. Uncontrolled liver disease

### REQUIRED MEDICAL INFORMATION

1. Dx of Sickle Cell Disease (SCD), 2. Tx to prevent acute complications of sickle cell disease, 3. Previous use, concurrent use, or inability to use generic hydroxyurea, 4. Reauthorization: reduction in the number of acute complications (i.e blood transfusions, sickle cell crisis, hospitalizations) of sickle cell disease since initiating therapy

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A

## **ENDOTHELIN ANTAGONISTS**

---

### **MEDICATION(S)**

ADEMPAS, AMBRISENTAN, BOSENTAN 125 MG TAB, BOSENTAN 62.5 MG TAB, OPSUMIT

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. PAH: a) Exclusion of all secondary causes of pulmonary hypertension b) Must be dx with PAH with WHO class II, III, or IV c) For Opsumit or Adempas: Previous trial of ambrisentan or bosentan 2. CTPH (for Adempas): a) Pt has failed endarterectomy OR b) Pt considered inoperable for pulmonary endarterectomy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ENSPRYNG**

---

### **MEDICATION(S)**

ENSPRYNG

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Use of another biologic tx for NMOSD

### **REQUIRED MEDICAL INFORMATION**

1. Dx of neuromyelitis optica spectrum disorder (NMOSD) with one of the following:

A. Idiopathic single or recurrent events of longitudinally extensive myelitis (3 or more vertebral segment spinal cord MRI lesion)

B. Optic neuritis, single, recurrent or simultaneous bilateral

2. Positive for anti-aquaporin-4 (AQP4) antibody,

3. Pt has tried and failed Uplizna or rituximab

4. Reauthorization: A. Patient is continuing to receive benefit from treatment

5. Chart notes required

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: Plan year Reauth: Plan Year

**OTHER CRITERIA**

N/A



## **ENTYVIO SQ**

---

### **MEDICATION(S)**

ENTYVIO PEN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of:
  - a. moderate to severe Crohn's disease
  - b. moderate to severe Ulcerative Colitis (UC)
2. For SQ, is patient established on Entyvio IV and changing to Entyvio SQ

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Gastroenterologist

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **EPCLUSA**

---

### **MEDICATION(S)**

SOFOSBUVIR-VELPATASVIR

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Chart notes documenting genotype,

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

12 Weeks

### **OTHER CRITERIA**

N/A

## **EPIDIOLEX**

---

### **MEDICATION(S)**

EPIDIOLEX

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

Previous use of two alternative antiepileptic medications and used in combination with another antiepileptic

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ERIVEDGE**

---

### **MEDICATION(S)**

ERIVEDGE, ODOMZO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has recurring lesions after radiation therapy OR radiation therapy is contraindicated or inappropriate, 2. Pt has recurring lesions after surgical excision OR surgery is contraindicated or inappropriate

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ERLEADA**

---

### **MEDICATION(S)**

ERLEADA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1)Diagnosis of metastatic castration sensitive prostate cancer (mCSPC) a)Patient has tried and failed abiraterone 250mg tablets AND Xtandi 2)Diagnosis of non-metastatic castration-resistant prostate cancer (nmCRPC) a)Patient has PSA doubling time of less than 10 months b)Patient has failed Xtandi

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ESBRIET**

---

### **MEDICATION(S)**

PIRFENIDONE

**PENDING CMS APPROVAL**

## **ETHACRYNIC ACID**

---

### **MEDICATION(S)**

ETHACRYNIC ACID

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt needs diuresis for any of the following: a. Edema associated with congestive heart failure, b. Edema associated with cirrhosis of the liver, c. Edema associates with renal disease. d. Short-term management of ascites due to malignancy, idiopathic edema, or lymphedema. 2. Pt has previous trial and failure on a loop diuretic or thiazide diuretic

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **EULEXIN**

---

### **MEDICATION(S)**

EULEXIN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Severe hepatic impairment

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of:
  - a. Locally confined Stage B2-C carcinoma of the prostate
  - b. Stage D2 metastatic carcinoma of the prostate
2. Will be used in combination with LHRH-agonist

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **EVENITY**

---

### **MEDICATION(S)**

EVENITY

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Duration not to exceed 24 months of cumulative treatment between all anabolic agents (Forteo, Tymlos, Evenity) unless the patient has returned to high risk or remains at high risk for fracture
2. Duration not to exceed 12 months with Evenity
3. Myocardial infarction or stroke in the last 12 months

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of osteoporosis or osteopenia
2. At least one of the following:
  - A. T-score worse than -3.5
  - B. T-score from -2.5 to -3.5 and at least one of the following:
    - i. History of multiple or recent fragility fracture
    - ii. T/f of oral or IV bisphosphonate or Prolia
  - C. T-score from -1.0 to -2.5 and BOTH of the following:
    - i. History of fragility fracture OR FRAX score of greater than 20% for major fracture or greater than 3% for hip fracture

ii. T/f of oral or IV bisphosphonate or Prolia

**AGE RESTRICTION**

N/A

**PRESCRIBER RESTRICTION**

N/A

**COVERAGE DURATION**

Plan year

**OTHER CRITERIA**

BvD Determination

## **FARESTON**

---

### **MEDICATION(S)**

TOREMIFENE CITRATE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Pt has congenital or acquired QT prolongation, 2. Pt has uncorrected hypokalemia, 3. Pt has uncorrected hypomagnesemia

### **REQUIRED MEDICAL INFORMATION**

1. Pt has previous trial and failure or contraindication to tamoxifen therapy, 2. Pt has previous trial and failure or contraindication to aromatase inhibitor therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **FASENRA**

---

### **MEDICATION(S)**

FASENRA, FASENRA PEN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Concurrent therapy with another biologic medication

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Severe Asthma      a. Patient has tried at least 3 months on any of the following combinations:  
    i. ICS/LABA      ii. ICS/LTRA      iii. ICS/LAMA      iv. ICS/LABA/LAMA (Trelegy)      b.

Two exacerbations requiring the use of oral corticosteroids in the previous 12 months OR one exacerbation that led to a hospitalization in the previous 12 months      c. Peripheral blood eosinophil level greater than 150 cells/mcL 2. Dx of eosinophilic granulomatosis with polyangiitis (EGPA)      a.

Patient must have TWO of the following disease characteristics      i. Biopsy showing histopathological evidence of eosinophilic vasculitis, perivascular eosinophilic infiltration, or eosinophil-rich granulomatous inflammation      ii. Neuropathy      iii. Pulmonary infiltrates, non-fixed  
    iv. Sino-nasal abnormality      v. Cardiomyopathy (established by echocardiography or MRI)  
    vi. Glomerulonephritis (hematuria, red cell casts, proteinuria)      vii. Alveolar hemorrhage  
    viii. Palpable purpura      ix. Anti-neutrophil cytoplasmic anti-body (ANCA) positive      c. Patient

has a history of eosinophil level greater than 100 cells/mcL or blood eosinophil level greater than 10%  
    d. Patient has a history of relapse or refractory disease despite current use of oral glucocorticoids unless contraindicated or not tolerated

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

For EGPA: allergist, immunologist, rheumatologist, or pulmonologist

### **COVERAGE DURATION**

Plan Year

## **OTHER CRITERIA**

BvsD Determination

## **FENTANYL PATCH**

---

### **MEDICATION(S)**

FENTANYL 100 MCG/HR PATCH 72HR, FENTANYL 12 MCG/HR PATCH 72HR, FENTANYL 25 MCG/HR PATCH 72HR, FENTANYL 50 MCG/HR PATCH 72HR, FENTANYL 75 MCG/HR PATCH 72HR

### **PENDING CMS APPROVAL**

## **FILSUVEZ**

---

### **MEDICATION(S)**

FILSUVEZ

**PENDING CMS APPROVAL**

## **FINTEPLA**

---

### **MEDICATION(S)**

FINTEPLA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Dravet Syndrome or Lennox Gaustaut Syndrome 2. Previous use of two of topiramate, valproic acid, or clobazam

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



# **FIRAZYR**

---

## **MEDICATION(S)**

ICATIBANT ACETATE

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

N/A

## **REQUIRED MEDICAL INFORMATION**

1. For HAE type I and II and acquired angioedema:

a. Dx has been verified by low C1-INH and/or low C1-INH function levels

2. For HAE with normal C1-INH:

a. Pt has a history of recurrent angioedema in the absence of urticaria and no use of medications known to cause angioedema

b. The pt has documented normal or near normal C4, C1-INH antigen, and C1-INH function

c. The patient has ONE of the following:

i. documentation showing a mutation associated with the disease OR

ii. a positive family history of recurrent angioedema and documented lack of efficacy of high-dose antihistamine therapy for at least 1 month or an interval expected to be associated with 3 or more attacks of angioedema (whichever is longer)

## **AGE RESTRICTION**

N/A

## **PRESCRIBER RESTRICTION**

N/A

**COVERAGE DURATION**

1. Initial: 6 months, 2. Reauth: Plan Year

**OTHER CRITERIA**

N/A

## **FIRDAPSE**

---

### **MEDICATION(S)**

FIRDAPSE

**PENDING CMS APPROVAL**

## **FOTIVDA**

---

### **MEDICATION(S)**

FOTIVDA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of relapsed or refractory advanced renal cell carcinoma (RCC) 2. Previous failure of two systemic therapies

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **FRUZAQLA**

---

### **MEDICATION(S)**

FRUZAQLA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of metastatic colorectal cancer (mCRC) 2. Member has previously been treated with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy, an anti-VEGF therapy, and, if RAS wild-type and medically appropriate, an anti-EGFR therapy 3. Member has failed, contraindication, or intolerance to Lonsurf with or without bevacizumab

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **GATTEX**

---

### **MEDICATION(S)**

GATTEX

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of short bowel syndrome
2. The patient is currently receiving parenteral nutrition/intravenous fluids at least 3 days per week
3. Dosing within FDA-approved dosage
3. Member has documented treatment failure, intolerance, or contraindication to octreotide therapy (octreotide treatment failure defined as lack of reduction in stool output within the first 4 weeks of treatment)
4. For adults, patient has had a colonoscopy in the last year and has had polyps removed if identified
5. For pediatric patients, a fecal occult blood test was performed and there was no new or unexplained blood in the stool OR a colonoscopy/sigmoidoscopy and upper GI endoscopy was performed if new or unexplained blood was identified in a fecal occult blood test
6. Reauthorization: Member has documented reduction in parental nutrition or intravenous support

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Gastroenterologist

### **COVERAGE DURATION**

6 months

### **OTHER CRITERIA**

N/A



## **GAVRETO**

---

### **MEDICATION(S)**

GAVRETO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of non-smallcell lung cancer (NSCLC) A. Metastatic NSCLC identified as rearranged during transfection (RET) fusion-positive 2. Dx of thyroid cancer A. Advanced or metastatic RET fusion-positive thyroid cancer refractory radioactive iodine (if appropriate) requiring systemic therapy

### **AGE RESTRICTION**

12 and older

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **GILOTRIF**

---

### **MEDICATION(S)**

GILOTRIF

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of metastatic non-small cell lung cancer (NSCLC) a. The tumor has an epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 (L858R) substitution mutation 2. Dx of metastatic squamous non-small cell lung cancer (NSCLC) a. Progression after platinum-based chemotherapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **GLEOSTINE**

---

### **MEDICATION(S)**

GLEOSTINE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of primary and metastatic brain tumors following appropriate surgical and/or radiotherapeutic procedures 2. Dx of Hodgkin's lymphoma A. Disease has progressed following initial chemotherapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **GLP 1**

---

### **MEDICATION(S)**

MOUNJARO, TRULICITY

### **PENDING CMS APPROVAL**

## **GOMEKLI**

---

### **MEDICATION(S)**

GOMEKLI

**PENDING CMS APPROVAL**

## **GROWTH HORMONE**

---

### **MEDICATION(S)**

GENOTROPIN, GENOTROPIN MINIQUEICK, OMNITROPE

### **PENDING CMS APPROVAL**

## **HAEGARDA**

---

### **MEDICATION(S)**

HAEGARDA, ORLADEYO

**PENDING CMS APPROVAL**

## **HARVONI**

---

### **MEDICATION(S)**

LEDIPASVIR-SOFOSBUVIR

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Chart notes showing genotype

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Maximum 24 weeks

### **OTHER CRITERIA**

N/A

## **HEMADY**

---

### **MEDICATION(S)**

HEMADY

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of Multiple Myeloma (MM) 2. Previous use of generic dexamethasone tablets

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## HEPATITIS C

---

### MEDICATION(S)

PEGASYS

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. For Hep B dx: a. Pre-treatment HBV DNA levels are greater than 20,000 IU/ml, b. Must be used as monotherapy 2. Hep C dx

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

48 weeks

### OTHER CRITERIA

N/A

## **HERNEXEOS**

---

### **MEDICATION(S)**

HERNEXEOS

### **PENDING CMS APPROVAL**

## **HETLIOZ**

---

### **MEDICATION(S)**

HETLIOZ LQ, TASIMELTEON

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For Non-24 Hour Sleep-Wake Disorder: a. Pt is totally blind without light perception 2. For Nighttime Sleep Disturbances in Smith-Magenis Syndrome (SMS) a. Diagnosis been confirmed with genetic testing

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Sleep specialist or Neurologist

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **HUMIRA**

---

### **MEDICATION(S)**

AMJEVITA 20 MG/0.2ML SOLN PRSYR, AMJEVITA 40 MG/0.4ML SOLN A-INJ, AMJEVITA 40 MG/0.4ML SOLN PRSYR, AMJEVITA 80 MG/0.8ML SOLN A-INJ, AMJEVITA-PED 15KG TO <30KG, HADLIMA, HADLIMA PUSHTOUCH

### **PENDING CMS APPROVAL**

## **HYFTOR**

---

### **MEDICATION(S)**

HYFTOR

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of facial angiofibroma associated with tuberous sclerosis 2. Member's facial angiofibroma cause functional impairment or symptoms such as bleeding or pain

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

12 weeks No reauthorization

### **OTHER CRITERIA**

N/A

## **IBRANCE**

---

### **MEDICATION(S)**

IBRANCE

**PENDING CMS APPROVAL**

## **IBTROZI**

---

### **MEDICATION(S)**

IBTROZI

**PENDING CMS APPROVAL**

## ICLUSIG

---

### MEDICATION(S)

ICLUSIG

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Chronic Phase CML dx a. Resistance or intolerance to at least two prior kinase inhibitors. 2. Accelerated or Blast phase CML OR Ph+ ALL a. No other TKI is indicated 3. T315I-positive CML or T315I-positive ALL 4. Newly diagnosed Ph+ ALL

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A



## **IDHIFA**

---

### **MEDICATION(S)**

IDHIFA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Cancer has an isocitrate dehydrogenase-2 (IDH2) mutation as detected by a FDA-approved test,
2. Pt has relapsed or is refractory to one or more prior anticancer regimens

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **IMBRUVICA**

---

### **MEDICATION(S)**

IMBRUVICA 140 MG CAP, IMBRUVICA 140 MG TAB, IMBRUVICA 280 MG TAB, IMBRUVICA 420 MG TAB, IMBRUVICA 70 MG CAP, IMBRUVICA 70 MG/ML SUSPENSION

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of CLL/SLL 2. Dx of CLL/SLL w/ 17p deletion 3. Dx of Waldenstrom's Macroglobinemia 4. Dx of cGVHD: a. Member has failed at least one prior systemic therapy b. Prescribing physician is a specialist in transplant

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **IMKELDI**

---

### **MEDICATION(S)**

IMKELDI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt is unable to swallow tablets

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **IMPAVIDO**

---

### **MEDICATION(S)**

IMPAVIDO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of leishmaniasis (visceral, cutaneous, or mucosal) confirmed by parasite in a clinical specimen 2. Previous use of ketoconazole, fluconazole, paromomycin, or amphotericin 3. Weight 30 kg or greater

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

28 days

### **OTHER CRITERIA**

N/A

## **INCRELEX**

---

### **MEDICATION(S)**

INCRELEX

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Height standard deviation score of less than -3 based on age and gender, 2. Basal IGF-1 standard deviation score of less than -3 based on age and gender, 3. Normal or elevated growth hormone levels, 4. Pt must have open epiphyses, 5. Gh stimulation test of greater than 10 mcg/L.

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## INLYTA

---

### MEDICATION(S)

INLYTA

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Dx of renal cell carcinoma 2. For first line treatment in combination with pembrolizumab (Keytruda) or avelumab (Bavencio) 3. Monotherapy as a second line treatment after previous use of one prior systemic therapy

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A

## **INQOVI**

---

### **MEDICATION(S)**

INQOVI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of myelodysplastic syndrome

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **INREBIC**

---

### **MEDICATION(S)**

INREBIC

**PENDING CMS APPROVAL**



# **IRESSA**

---

## **MEDICATION(S)**

GEFITINIB

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

N/A

## **REQUIRED MEDICAL INFORMATION**

1. Pt has ONE of the following: a. EGFR exon 19 deletion, b. EGFR exon 21 deletion, 2. Gefitinib is used as first-line therapy

## **AGE RESTRICTION**

N/A

## **PRESCRIBER RESTRICTION**

N/A

## **COVERAGE DURATION**

Plan Year

## **OTHER CRITERIA**

N/A

## **ISTURISA**

---

### **MEDICATION(S)**

ISTURISA 1 MG TAB, ISTURISA 5 MG TAB

### **PENDING CMS APPROVAL**

## **ITOVEBI**

---

### **MEDICATION(S)**

ITOVEBI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Disease progression on or following other PI3k inhibitors

### **REQUIRED MEDICAL INFORMATION**

1. Pt must have advanced or metastatic HR-positive, HER2-negative breast cancer with documented PIK3CA mutation via specified testing 2. Pt must have disease progression following at least one line of endocrine therapy in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy 3. Itovebi will be given in combination with Ibrance (palbociclib) and fulvestrant as first-line therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **IVIG**

---

### **MEDICATION(S)**

BIVIGAM, GAMMAGARD, GAMMAGARD S/D LESS IGA, GAMMAKED 1 GM/10ML SOLUTION, GAMMAPLEX, GAMUNEX-C 1 GM/10ML SOLUTION, OCTAGAM 1 GM/20ML SOLUTION, OCTAGAM 10 GM/100ML SOLUTION, OCTAGAM 10 GM/200ML SOLUTION, OCTAGAM 2 GM/20ML SOLUTION, OCTAGAM 2.5 GM/50ML SOLUTION, OCTAGAM 25 GM/500ML SOLUTION, OCTAGAM 30 GM/300ML SOLUTION, OCTAGAM 5 GM/100ML SOLUTION, PRIVIGEN

### **PA INDICATION INDICATOR**

3 - All Medically-Accepted Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1)For all non-preferred IVIG products (ie Bivigam, Gammaked, Octagam, etc) a)Must try TWO of the following: Gammagard, Gamunex, Hizentra, and Privigen OR b)Documented medical reason why preferred agents cannot be used

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

BvD Determination

## **IWILFIN**

---

### **MEDICATION(S)**

IWILFIN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of high risk neuroblastoma (HRNB) in patients who have demonstrated at least a partial response to prior multiagent, multimodality therapy including anti-GD2 immunotherapy 2. Maximum duration of 2 years of therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## JADENU

---

### MEDICATION(S)

DEFERASIROX 180 MG PACKET, DEFERASIROX 250 MG TAB SOL, DEFERASIROX 360 MG PACKET, DEFERASIROX 500 MG TAB SOL, DEFERASIROX 90 MG PACKET, DEFERASIROX GRANULES

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. For blood transfusion dx: a. Pt's serum ferritin is greater than 1000 mcg/L, b. Pt has failed subcutaneous deferoxamine through an inability to achieve desired goals of therapy or been intolerant to therapy, 2. For non-transfusion-dependent thalassemia dx: a. Pt's liver iron concentration is at least 5 mg Fe per gram of dry weight, b. Pt's serum ferritin is greater than 300 mcg/L, c. Pt has failed subcutaneous deferoxamine through an inability to achieve desired goals of therapy or been intolerant to therapy

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

Part B before Part D step therapy

## **JAKAFI**

---

### **MEDICATION(S)**

JAKAFI

**PENDING CMS APPROVAL**

## JAYPIRCA

---

### MEDICATION(S)

JAYPIRCA

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Diagnosis of relapsed/refractory MCL    a. Previous treatment with at least two lines of systemic therapy, including a BTK inhibitor  
2. Diagnosis of chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL)    a. Previous treatment with at least two prior lines of therapy including a BTK inhibitor and a BCL-2 inhibitor

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A



## JYNARQUE

---

### MEDICATION(S)

TOLVAPTAN 15 MG TAB THPK, TOLVAPTAN 30 & 15 MG TAB THPK, TOLVAPTAN 45 & 15 MG TAB THPK, TOLVAPTAN 60 & 30 MG TAB THPK, TOLVAPTAN 90 & 30 MG TAB THPK

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

eGFR less than 25ml/min

### REQUIRED MEDICAL INFORMATION

1. Diagnosis of autosomal dominant polycystic kidney disease    a) Diagnosis has been confirmed by radiology

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

Endocrinologist, nephrologist, cardiologist and hepatologist.

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A

## **KALYDECO**

---

### **MEDICATION(S)**

KALYDECO

**PENDING CMS APPROVAL**

## **KERENDIA**

---

### **MEDICATION(S)**

KERENDIA

**PENDING CMS APPROVAL**

## **KEVEYIS**

---

### **MEDICATION(S)**

KEVEYIS

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Hypersensitivity to dichlorphenamide or other sulfonamides 2. Use in combination with high-dose aspirin 3. Severe pulmonary disease 4. Hepatic insufficiency (e.g Child-Pugh B or C)

### **REQUIRED MEDICAL INFORMATION**

1. Dx of primary hyperkalemic periodic paralysis, primary hypokalemic periodic paralysis, or related variants. 2. Dx confirmed by genetic testing, provocative testing, electromyography, or muscle biopsy 3. Previous use or contraindication to acetazolamide 4. Reauthorization: a. improvement in baseline symptoms (e.g. number of attacks per week or month, severity of attacks, duration of attacks, short-form 36 assessment, etc.)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: 6 months Reauthorization: Plan Year

### **OTHER CRITERIA**

N/A

## **KISQALI**

---

### **MEDICATION(S)**

KISQALI (200 MG DOSE), KISQALI (400 MG DOSE), KISQALI (600 MG DOSE), KISQALI FEMARA (200 MG DOSE), KISQALI FEMARA (400 MG DOSE), KISQALI FEMARA (600 MG DOSE)

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer, must meet ONE of the following: a. Pt will receive an aromatase inhibitor in combination with Kisqali as initial endocrine based therapy for advanced or metastatic disease, b. Pt will receive fluvestrant in combination with Kisqali 2. Dx of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative stage II or stage III early breast cancer at high risk of recurrence a. Pt will receive an aromatase inhibitor in combination with Kisqali

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **KORLYM**

---

### **MEDICATION(S)**

MIFEPRISTONE 300 MG TAB

**PENDING CMS APPROVAL**

## **KOSELUGO**

---

### **MEDICATION(S)**

KOSELUGO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Patient has symptomatic, inoperable plexiform neurofibromas (PN)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## KYNMOBI

---

### MEDICATION(S)

APOMORPHINE HCL

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Dx of Parkinson's disease A. Experiencing off episodes 2. Currently taking an oral formulation of carbidopa/levodopa or there is documentation of an inability to take an oral formulation of carbidopa/levodopa 3. Previous use of an immediate release or oral disintegrating carbidopa/levodopa as a rescue for off episodes 4. Previous use of at least one of: a. COMT inhibitor (tolcapone, entacapone), b. Dopamine agonist (ropinirole, pramipexole), c. MAO-B inhibitor (selegiline, rasagiline, safinamide) 5. Prescribed in combination with antiemetic therapy (Not a 5HT3)

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A



## **LAMPIT**

---

### **MEDICATION(S)**

LAMPIT

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Chagas disease (*T. cruzi*)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

3 months

### **OTHER CRITERIA**

N/A

## **LAZCLUZE**

---

### **MEDICATION(S)**

LAZCLUZE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Prior systemic treatment for locally advanced Stage III or metastatic Stage IV disease 2. Symptomatic or previously treated unstable brain metastases

### **REQUIRED MEDICAL INFORMATION**

1. Initial A. Patient must have confirmed NSCLC that is metastatic or unresectable with EGFR exon 19 deletions or 21 L858R substitution mutations. B. Patient must receive anticoagulant prophylaxis to prevent VTE events for the first 4 months of treatment. C. Patients must have an ECOG of 0 or 1 D. Clinical rationale why Tagrisso with /without chemotherapy is not appropriate for use in the patient. 2. Reauthorization A. Tumor assessment does not show new growth, activity or mutations

### **AGE RESTRICTION**

18 years of age or older

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

6 months

### **OTHER CRITERIA**

N/A

## **LENVIMA**

---

### **MEDICATION(S)**

LENVIMA (10 MG DAILY DOSE), LENVIMA (12 MG DAILY DOSE), LENVIMA (14 MG DAILY DOSE), LENVIMA (18 MG DAILY DOSE), LENVIMA (20 MG DAILY DOSE), LENVIMA (24 MG DAILY DOSE), LENVIMA (4 MG DAILY DOSE), LENVIMA (8 MG DAILY DOSE)

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Renal cell carcinoma dx: a. Will be used in combination with everolimus OR pembrolizumab 2. Thyroid cancer dx: a. Tumor is refractory to treatment with radioactive iodine, b. Used as monotherapy 3. Unresectable hepatocellular carcinoma dx: 4. Endometrial carcinoma dx: a. Will be used in combination with pembrolizumab (Keytruda) b. Does not have microsatellite instability-high or mismatch repair deficiency c. Pt has previously been treated with systemic therapy d. Pt is not a candidate for surgery or radiation

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **LEUKERAN**

---

### **MEDICATION(S)**

LEUKERAN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of:
  - a. chronic lymphatic (lymphocytic) leukemia
  - b. malignant lymphomas including lymphosarcoma,
  - c. giant follicular lymphoma
  - d. Hodgkins disease

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

# LEUKINE

---

## MEDICATION(S)

LEUKINE

## PA INDICATION INDICATOR

1 - All FDA-Approved Indications

## OFF LABEL USES

N/A

## EXCLUSION CRITERIA

N/A

## REQUIRED MEDICAL INFORMATION

1. Dx of: a. To shorten time to neutrophil recovery and to reduce the incidence of severe and life-threatening infections and infections resulting in death following induction chemotherapy in adult patients 55 years and older with acute myeloid leukemia (AML). i. Pt has trial on BOTH of the following: a. Fulphila, b. Udenyca b, For the mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation. c. For the acceleration of myeloid reconstitution following autologous bone marrow or peripheral blood progenitor cell transplantation. d. For the acceleration of myeloid reconstitution following allogeneic bone marrow transplantation. e. For treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic bone marrow transplantation. f. To increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).

## AGE RESTRICTION

N/A

## PRESCRIBER RESTRICTION

N/A

## COVERAGE DURATION

Plan Year

## OTHER CRITERIA

BvD Determination



## **LIVTENCITY**

---

### **MEDICATION(S)**

LIVTENCITY

**PENDING CMS APPROVAL**

## **LONSURF**

---

### **MEDICATION(S)**

LONSURF

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

A. Metastatic colorectal cancer 1. Pt has previous therapy on the following: a. A fluoropyrimidine, b. Oxaliplatin, c. Irinotecan, d. Bevacizumab, 2. If cancer is KRAS wild type, pt has received previous therapy with anti-EGFR therapy B. Metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least two prior lines of chemotherapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **LORBRENA**

---

### **MEDICATION(S)**

LORBRENA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1) Dx of metastatic non-small cell lung cancer (NSCLC) a) Tumors are anaplastic lymphoma kinase (ALK)-positive 2) Patient has failed/intolerance/contraindication to Alecensa

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **LUCEMYRA**

---

### **MEDICATION(S)**

LOFEXIDINE HCL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Mitigation of opioid withdrawal symptoms 2. Provider submitted documentation that the patient has been counseled on the risks of taking lofexidine with alcohol, benzodiazepines, and/or barbituates 3. Patient has failed clonidine as part of this opioid discontinuation attempt

### **AGE RESTRICTION**

18 years of age or older

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

14 Days

### **OTHER CRITERIA**

N/A

## **LUMAKRAS**

---

### **MEDICATION(S)**

KRAZATI, LUMAKRAS

**PENDING CMS APPROVAL**

## **LUPKYNIS**

---

### **MEDICATION(S)**

LUPKYNIS

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of lupus nephritis 2. eGFR greater than 45 mL/min/1.73m<sup>2</sup>

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **LYNPARZA**

---

### **MEDICATION(S)**

LYNPARZA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Combination therapy

### **REQUIRED MEDICAL INFORMATION**

1. Ovarian cancer, advanced (BRCA-mutated): a. First-line maintenance therapy for gBRCAm or sBRCAm advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in adult patients with complete or partial response to first-line platinum-based chemotherapy, OR b. First-line maintenance treatment (in combination with bevacizumab) of advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in adults who are in complete or partial response to first-line, platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency (HRD)-positive status, 2. Recurrent ovarian cancer dx: a. Maintenance treatment of recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer in adults who are in complete or partial response to platinum-based chemotherapy. 3. Breast Cancer (BRCA-mutated, HER2-negative) dx: a. For Metastatic Breast Cancer: Pt has previous trial of chemotherapy in neoadjuvant, adjuvant, or metastatic setting i. If HR positive: pt has t/f endocrine therapy or endocrine therapy is inappropriate for pt b. For High Risk Early Breast Cancer: Pt has previous trial of chemotherapy in neoadjuvant, adjuvant, or metastatic setting 4. Pancreatic cancer (BRCA-mutated): a. disease has not progressed on at least 16 weeks of a first-line, platinum-based chemotherapy regimen 5. Prostate cancer (mCRPC): a. Homologous recombination repair (HRR) gene-mutated metastatic castration-resistant prostate cancer in adults who have progressed following prior enzalutamide or abiraterone treatment b. BRCA-mutated (BRCAm) metastatic castration resistant prostate cancer (mCRPC) in combination with abiraterone and prednisone or prednisolone

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

**COVERAGE DURATION**

Plan Year

**OTHER CRITERIA**

N/A

## **LYTGOBI**

---

### **MEDICATION(S)**

LYTGOBI (12 MG DAILY DOSE), LYTGOBI (16 MG DAILY DOSE), LYTGOBI (20 MG DAILY DOSE)

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of metastatic cholangiocarcinoma 2. Previous treatment for metastatic cholangiocarcinoma with at least 1 line of systemic therapy 3. Fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **MAVYRET**

---

### **MEDICATION(S)**

MAVYRET

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

Confirmation of genotype

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

16 Weeks

### **OTHER CRITERIA**

N/A



## **MEKINIST**

---

### **MEDICATION(S)**

MEKINIST

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Will be used as monotherapy for:

a. Dx of unresectable or metastatic melanoma with BRAF V600E or V600K mutations in BRAF-inhibitor treatment-naïve patients

2. Will be used in combination with dabrafenib for:

a. Dx of unresectable or metastatic melanoma with BRAF V600E or V600K mutations

b. The adjuvant treatment of patients with dx of melanoma with BRAF V600E or V600K mutations and involvement of lymph node(s), following complete resection.

c. Dx of patients with metastatic non-small cell lung cancer (NSCLC) with BRAF V600E mutation

d. Dx of patients with locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and with no satisfactory locoregional treatment options.

e. Dx of unresectable or metastatic solid tumors with BRAF V600E mutation who have progressed following prior treatment and have no satisfactory alternative treatment options.

f. Dx of low-grade glioma (LGG) with a BRAF V600E mutation who require systemic therapy.

3. For oral solution: member has inability to swallow tablets

**AGE RESTRICTION**

N/A

**PRESCRIBER RESTRICTION**

N/A

**COVERAGE DURATION**

Plan Year

**OTHER CRITERIA**

N/A

## **MIGRANAL**

---

### **MEDICATION(S)**

DIHYDROERGOTAMINE MESYLATE 4 MG/ML SOLUTION

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt experiences at least 2 migraines per month, 2. Pt has trial or contraindication to at least TWO of the following: a. Sumatriptan, b. Rizatriptan, c. Zolmitriptan, d. Naratriptan

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **MODEYSO**

---

### **MEDICATION(S)**

MODEYSO

**PENDING CMS APPROVAL**

## **MULPLETA**

---

### **MEDICATION(S)**

MULPLETA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Platelet count less than 50,000 2. Pt is scheduled for a procedure where there is a bleeding risk 3. Mulpleta will be used for 7 days starting 8 to 14 days prior to the procedure and discontinued 2 to 8 days prior to the procedure

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **MYALEPT**

---

### **MEDICATION(S)**

MYALEPT

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has baseline leptin levels of less than 8 ng/mL for males OR less than 12 ng/mL for females, 2. Pt has ONE of the following: a. Diagnosis of diabetes and is being treated with Metformin AND at least one other antidiabetic agent, b. Diagnosis of hypertriglyceridemia and is being treated with at least ONE antihyperlipidemic agent, 3. Reauth: a. Pt has been screened for the presence of anti-metrelptin antibodies, b. If presence of anti-metrelptin antibodies, pt must still be receiving benefit from Myalept therapy, c. Pt shows improvement in hemoglobin A1c OR fasting triglyceride level

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: 6 months, Reauth: Plan Year

### **OTHER CRITERIA**

N/A

## **NARCOLEPSY**

---

### **MEDICATION(S)**

WAKIX

### **PA INDICATION INDICATOR**

2 - Some FDA-Approved Indications Only

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Patients with severe hepatic impairment

### **REQUIRED MEDICAL INFORMATION**

1. For narcolepsy with cataplexy (please note Wakix is only covered for the narcolepsy with cataplexy indication) a. Patient has been diagnosed by a board certified sleep, pulmonology, or neurology specialist b. Patient exhibits symptoms of cataplexy 2. Reauth: a. For narcolepsy with cataplexy: i. Decrease in cataplexy episodes

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Board certified in sleep, pulmonology, or neurology

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **NERLYNX**

---

### **MEDICATION(S)**

NERLYNX

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Breast cancer dx: a. Tx of early stage HER2-positive breast cancer following adjuvant trastuzumab-based therapy OR b. Tx of HER2-positive breast cancer, in combination with capecitabine, in patients who have received 2 or more prior regimens for metastatic disease

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **NEULASTA**

---

### **MEDICATION(S)**

FYLNETRA, NEULASTA, NYVEPRIA, STIMUFEND, ZIEXTENZO

### **PENDING CMS APPROVAL**

## **NEXAVAR**

---

### **MEDICATION(S)**

SORAFENIB TOSYLATE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Unresectable hepatocellular carcinoma
2. Dx of Advanced renal cell carcinoma
3. Dx of Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma (DTC) refractory to radioactive iodine treatment

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **NEXLETOL**

---

### **MEDICATION(S)**

NEXLETOL

**PENDING CMS APPROVAL**

## **NEXLIZET**

---

### **MEDICATION(S)**

NEXLIZET

**PENDING CMS APPROVAL**

## **NICOTROL**

---

### **MEDICATION(S)**

NICOTROL NS

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has tried varenicline or bupropion OR has a contraindication to the use of varenicline or bupropion (established seizure disorder, concurrent anorexia or bulimia, concurrent use of MAOIs, etc)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **NINLARO**

---

### **MEDICATION(S)**

NINLARO

### **PA INDICATION INDICATOR**

4 - All FDA-Approved Indications, Some Medically-Accepted Indications

### **OFF LABEL USES**

Autologous Stem Cell Transplant

### **EXCLUSION CRITERIA**

1. Pt is refractory to lenalidomide or proteasome inhibitor therapy

### **REQUIRED MEDICAL INFORMATION**

1. Multiple Myeloma

a. Combination with lenalidomide (Revlimid) and dexamethasone

b. Pt has previous trial on at least ONE other therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **NITROFURAN**

---

### **MEDICATION(S)**

NITROFURANTOIN 25 MG/5ML SUSPENSION, NITROFURANTOIN 50 MG/10ML SUSPENSION

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Unable to swallow nitrofurantoin capsules

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **NORTHERA**

---

### **MEDICATION(S)**

DROXIDOPA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has previous trial on BOTH of the following: a. Midorine, b. Fludrocortisone

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **NOXAFIL**

---

### **MEDICATION(S)**

NOXAFIL 300 MG PACKET

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For treatment of invasive aspergillus 2. For aspergillus or candida prophylaxis: a. Pt is at high risk of developing infections secondary to being severely immunocompromised, 3. For oropharyngeal candidiasis: a. Pt has previous failure on BOTH of the following: 1) Itraconazole, 2) Fluconazole

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

6 Months

### **OTHER CRITERIA**

N/A

## **NUBEQA**

---

### **MEDICATION(S)**

NUBEQA

**PENDING CMS APPROVAL**

## **NUPLAZID**

---

### **MEDICATION(S)**

NUPLAZID

**PENDING CMS APPROVAL**

## **NURTEC**

---

### **MEDICATION(S)**

NURTEC

### **PA INDICATION INDICATOR**

2 - Some FDA-Approved Indications Only

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For Acute Treatment of Migraines (Please note: Nurtec is only covered for acute treatment of migraines, prophylaxis dosing is not covered): a. Patient has tried 2 generic triptan medications OR has a contraindication to the use of triptans (e.g. established cardiovascular disease or cerebrovascular disease)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **OCTREOTIDE**

---

### **MEDICATION(S)**

OCTREOTIDE ACETATE 100 MCG/ML SOLUTION, OCTREOTIDE ACETATE 1000 MCG/ML SOLUTION, OCTREOTIDE ACETATE 200 MCG/ML SOLUTION, OCTREOTIDE ACETATE 50 MCG/ML SOLUTION, OCTREOTIDE ACETATE 500 MCG/ML SOLUTION

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For acromegaly: A. Has patient failed at least TWO of the following: i. Surgical resection, ii. Pituitary irradiation, iii. Bromocriptine, 2. Dx of metastatic carcinoid tumors: c. Dx of vasoactive intestinal peptide secreting tumors (VIPoma)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

BvsD determination

**OFEV**

---

**MEDICATION(S)**

OFEV

**PENDING CMS APPROVAL**

## **OGSIVEO**

---

### **MEDICATION(S)**

OGSIVEO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of desmoid tumor/aggressive fibromatosis with documentation of tumor progression 2.  
Contraindication, intolerance, or failure of sorafenib

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **OJEMDA**

---

### **MEDICATION(S)**

OJEMDA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Patient with tumors harboring additional activating molecular alteration(s) (e.g. IDH1/2 mutations, FGFR mutations, etc) or patients with known or suspected diagnosis of neurofibromatosis type 1 (NF1)

### **REQUIRED MEDICAL INFORMATION**

1. Dx of relapsed or refractory pediatric low-grade glioma (LGG) harboring an activating BRAF alteration based on local laboratory testing 2. At least one measurable lesion as defined by RANO 2010 criteria 3. Pt has received at least one line of prior systemic therapy and had documented evidence of radiographic progression 4. Pt has contraindication, intolerance, or failure of Tafenlar and Mekinist if BRAF V600 positive

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **OJJAARA**

---

### **MEDICATION(S)**

OJJAARA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of intermediate or high-risk myelofibrosis (MF), including primary MF or secondary MF (postpolycythemia vera (PV) and post-essential thrombocythemia (ET)) 2. Hemoglobin less than 10g/dL 3. Member has tried and failed or has intolerance to Jakafi

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## ONFI

---

### MEDICATION(S)

SYMPAZAN

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Pt has previous trail on at least TWO AED medications

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A

## **ONUREG**

---

### **MEDICATION(S)**

ONUREG

**PENDING CMS APPROVAL**

## **ORFADIN**

---

### **MEDICATION(S)**

NITISINONE, NITYR, ORFADIN 4 MG/ML SUSPENSION

**PENDING CMS APPROVAL**

## **ORGOVYX**

---

### **MEDICATION(S)**

ORGOVYX

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Previous treatment with another GnRH/LHR agonist/antagonist

### **REQUIRED MEDICAL INFORMATION**

1. Dx of castrate-sensitive metastatic prostate cancer 2. Dx of metastatic disease has been confirmed by bone scan, ultrasound, CT, MRI, or biopsy 3. Serum PSA is elevated 4. Contraindication or inability to take other GNRH/LHR agonist/antagonist medication due to one of: A. Short term (6 month) use in men at risk of toxicities from standard androgen deprivation therapy (ADT) B. Intermittent ADT in frail patients at risk of ADT toxicities C. Significant underlying cardiac risk factors

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ORIAHNN**

---

### **MEDICATION(S)**

MYFEMBREE, ORIAHNN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Maximum lifetime duration 24 months

### **REQUIRED MEDICAL INFORMATION**

1)For Oriahnn and Myfembree: Heavy menstrual bleeding associated with uterine leiomyomas (fibroids) 2)For Myfembree: Moderate to Severe Pain Associated with Endometriosis 3)Premenopausal 4)Previous use of a combination oral contraceptive 5)Previous use of a progestin-only contraceptive

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ORILISSA**

---

### **MEDICATION(S)**

ORILISSA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1) previous use of combination oral contraceptive 2) previous use of progestin-only contraceptive

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year Orilissa 150 mg, maximum lifetime duration 24 months 6 months Orilissa 200 mg

### **OTHER CRITERIA**

N/A

## **ORKAMBI**

---

### **MEDICATION(S)**

ORKAMBI

**PENDING CMS APPROVAL**



## **ORSERDU**

---

### **MEDICATION(S)**

ORSERDU

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Postmenopausal female or male with a diagnosis of advanced or metastatic ER+, HER2- breast cancer 2. Confirmation of ESR1-mutated breast cancer 3. The member has experienced disease progression following at least one line of endocrine therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

# **OSTEOPOROSIS**

---

## **MEDICATION(S)**

TERIPARATIDE, TYMLOS

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

1. Duration not to exceed 24 months of cumulative treatment between all anabolic agents (Forteo, Tymlos, Evenity) unless the patient has returned to high risk or remains at high risk for fracture

## **REQUIRED MEDICAL INFORMATION**

1. For teriparatide:

A. Member must fail Tymlos OR

B. Member has glucocorticoid-associated osteoporosis

2. Diagnosis of osteoporosis or osteopenia

3. At least one of the following:

A. T-score worse than -3.5

B. T-score from -2.5 to -3.5 and at least one of the following:

i. History of multiple or recent fragility fracture

ii. T/f of oral or IV bisphosphonate or Prolia

C. T-score from -1.0 to -2.5 and BOTH of the following:

i. History of fragility fracture OR FRAX score of greater than 20% for major fracture or greater than

3% for hip fracture

ii. T/f of oral or IV bisphosphonate or Prolia

**AGE RESTRICTION**

N/A

**PRESCRIBER RESTRICTION**

N/A

**COVERAGE DURATION**

Plan year

**OTHER CRITERIA**

N/A

## **PAH**

---

### **MEDICATION(S)**

SILDENAFIL CITRATE 10 MG/ML RECON SUSP, SILDENAFIL CITRATE 20 MG TAB, TADALAFIL (PAH), TADLIQ

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Initial: a. Pt is diagnosed with pulmonary arterial hypertension, confirmed by right heart catheterization b. Pt with positive vasoreactivity test: i. Pt has contraindications to or failed maximum tolerated doses of calcium channel blockers, c. For Tadliq and sildenafil oral suspension requests: Inability to swallow tablets 2. Reauth: a. Pt has been reassessed within the past 6 months

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **PALYNZIQ**

---

### **MEDICATION(S)**

PALYNZIQ

**PENDING CMS APPROVAL**

## **PANRETIN**

---

### **MEDICATION(S)**

PANRETIN

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Systemic anti-KS therapy is required (e.g., more than 10 new KS lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary KS, or symptomatic visceral involvement)

### **REQUIRED MEDICAL INFORMATION**

1. Dx of cutaneous lesions in patients with AIDS-related Kaposi's Sarcoma. 2. Reauthorization: Patient is stable on therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: 6 months Reauth: Plan Year

### **OTHER CRITERIA**

N/A

## PART D VS PART B

---

### MEDICATION(S)

ABILIFY ASIMTUFII, ABILIFY MAINTENA, ACETYLCYSTEINE 10 % SOLUTION, ACETYLCYSTEINE 20 % SOLUTION, ALBUTEROL SULFATE (2.5 MG/3ML) 0.083% NEBU SOLN, ALBUTEROL SULFATE (5 MG/ML) 0.5% NEBU SOLN, ALBUTEROL SULFATE 0.63 MG/3ML NEBU SOLN, ALBUTEROL SULFATE 1.25 MG/3ML NEBU SOLN, ALBUTEROL SULFATE 2.5 MG/0.5ML NEBU SOLN, APREPITANT, ARANESP (ALBUMIN FREE), ARFORMOTEROL TARTRATE, ARISTADA, ARISTADA INITIO, ASTAGRAF XL, AZATHIOPRINE, BUDESONIDE 0.25 MG/2ML SUSPENSION, BUDESONIDE 0.5 MG/2ML SUSPENSION, BUDESONIDE 1 MG/2ML SUSPENSION, CROMOLYN SODIUM 20 MG/2ML NEBU SOLN, CYCLOPHOSPHAMIDE 25 MG CAP, CYCLOPHOSPHAMIDE 25 MG TAB, CYCLOPHOSPHAMIDE 50 MG CAP, CYCLOPHOSPHAMIDE 50 MG TAB, CYCLOSPORINE 100 MG CAP, CYCLOSPORINE 25 MG CAP, CYCLOSPORINE MODIFIED, ELIGARD 22.5 MG KIT, ELIGARD 30 MG KIT, ELIGARD 7.5 MG KIT, ENGERIX-B, ENVARSUS XR, EPOGEN, ERZOFRI, EVEROLIMUS 0.25 MG TAB, EVEROLIMUS 0.5 MG TAB, EVEROLIMUS 0.75 MG TAB, EVEROLIMUS 1 MG TAB, FIRMAGON, FIRMAGON (240 MG DOSE), FLUPHENAZINE DECANOATE, FLUPHENAZINE HCL 2.5 MG/ML SOLUTION, FORMOTEROL FUMARATE, FULPHILA, GENGRAF 100 MG CAP, GENGRAF 25 MG CAP, GRANISETRON HCL 1 MG TAB, GRANIX 300 MCG/0.5ML SOLN PRSYR, GRANIX 300 MCG/ML SOLUTION, GRANIX 480 MCG/0.8ML SOLN PRSYR, HALOPERIDOL DECANOATE, HALOPERIDOL LACTATE 5 MG/ML SOLUTION, HEPLISAV-B, INVEGA HAFYERA, INVEGA SUSTENNA, INVEGA TRINZA, IPRATROPIUM BROMIDE 0.02 % SOLUTION, IPRATROPIUM-ALBUTEROL, JUBBONTI, LEUPROLIDE ACETATE (3 MONTH), LEVALBUTEROL HCL, LILETTA (52 MG), LUPRON DEPOT (1-MONTH), LUPRON DEPOT (3-MONTH), LUPRON DEPOT (4-MONTH), LUPRON DEPOT (6-MONTH), LUPRON DEPOT-PED (1-MONTH) 7.5 MG KIT, LUPRON DEPOT-PED (3-MONTH) 11.25 MG (PED) KIT, METHOTREXATE SODIUM 250 MG/10ML SOLUTION, METHOTREXATE SODIUM 50 MG/2ML SOLUTION, METHOTREXATE SODIUM (PF), MICAfungin SODIUM, MYCOPHENOLATE MOFETIL 200 MG/ML RECON SUSP, MYCOPHENOLATE MOFETIL 250 MG CAP, MYCOPHENOLATE MOFETIL 500 MG TAB, MYCOPHENOLATE SODIUM, MYCOPHENOLIC ACID, NIVESTYM, OLANZAPINE 10 MG RECON SOLN, ONDANSETRON 4 MG TAB DISP, ONDANSETRON 8 MG TAB DISP, ONDANSETRON HCL 4 MG TAB, ONDANSETRON HCL 4 MG/5ML SOLUTION, ONDANSETRON HCL 8 MG TAB, PENTAMIDINE ISETHIONATE, PERSERIS, PROGRAF 0.2 MG PACKET, PROGRAF 1 MG PACKET, PULMOZYME, RECOMBIVAX HB, RETACRIT, RISPERIDONE MICROSPHERES ER, SIROLIMUS, SODIUM CHLORIDE 0.9 % SOLUTION, STREPTOMYCIN SULFATE, TACROLIMUS 0.5 MG CAP, TACROLIMUS 1 MG CAP, TACROLIMUS 5 MG CAP, TESTOSTERONE CYPIONATE, TESTOSTERONE ENANTHATE, TRELSTAR MIXJECT, TWINRIX, UDENYCA, UZEDY, VARUBI (180 MG DOSE), WYOST,

## ZIPRASIDONE MESYLATE

### **DETAILS**

This drug may be covered under Medicare Part B or D depending on the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.



## **PCSK**

---

### **MEDICATION(S)**

REPATHA, REPATHA SURECLICK

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. The patient meets one of the following: A. The patient has tried one high-intensity statin(e.g. atorvastatin 40-80mg, rosuvastatin 20-40mg) or a maximally tolerated statin without achieving the LDL-C goal B. The patient experienced statin-related rhabdomyolysis with documented CK elevations greater than 10x ULN C. The patient experienced muscle-related symptoms such as myopathy or myalgia while on two separate trials of different statin therapy that both resolved upon discontinuation of statin therapy 2. AND meets ONE of the following: A. Dx of HoFH , untreated LDL-C greater than 500mg/dL or treated LDL-C greater than 300mg/dL i. AND cutaneous or tendon xanthoma before age 10 years, OR ii. Elevated LDL-C levels consistent with heterozygous FH in both parents B. Dx of HeFH or primary hyperlipidemia with fasting LDL-C of 190 mg/dL or greater on at least two separate dates at least 3 months apart i. AND LDL-C remains greater than 100 mg/dL despite treatment on medication therapy C. Dx of ASCVD consisting of MI, stroke, TIA, persistent intermittent claudication, coronary intervention revascularization or angina with proven ischemia i. AND LDL-C remains greater than 70 mg/dL despite treatment on medication therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

**OTHER CRITERIA**

N/A

## **PEMAZYRE**

---

### **MEDICATION(S)**

PEMAZYRE

**PENDING CMS APPROVAL**

## **PHENOXYBENZAMINE**

---

### **MEDICATION(S)**

METYROSINE, PHENOXYBENZAMINE HCL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Phenoxybenzamine will be used for short-term treatment of hypertension prior to surgical removal of a pheochromocytoma 2. For metyrosine, pt has failed phenoxybenzamine

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

3 mo

### **OTHER CRITERIA**

N/A

## **PIQRAY**

---

### **MEDICATION(S)**

PIQRAY (200 MG DAILY DOSE), PIQRAY (250 MG DAILY DOSE), PIQRAY (300 MG DAILY DOSE)

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, PIK3CA-mutated, advanced or metastatic breast cancer    a. PIK3CA-mutation positive    b. Receiving or previous use of an endocrine-based regimen

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **POMALYST**

---

### **MEDICATION(S)**

POMALYST

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Multiple myeloma dx: a. Pt has tried BOTH of the following: i. Revlimid, ii. bortezomib, b. Pt has demonstrated disease progression within 60 days of completion of prior therapy 2. Kaposi sarcoma dx: a. Experienced failure of highly active antiretroviral therapy (HAART) in patient with AIDS b. HIV-negative

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **PRETOMANID**

---

### **MEDICATION(S)**

PRETOMANID

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of tuberculosis 2. Used as part of an appropriate treatment regimen (e.g. bedaquiline and linezolid)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

6 months

### **OTHER CRITERIA**

N/A

## **PREVYMIS**

---

### **MEDICATION(S)**

PREVYMIS 120 MG PACKET, PREVYMIS 20 MG PACKET, PREVYMIS 240 MG TAB, PREVYMIS 480 MG TAB

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Pt has Child-Pugh class C hepatic impairment

### **REQUIRED MEDICAL INFORMATION**

1. For: CMV prophylaxis in HSCT recipients a. Pt is post allogenic hematopoietic stem cell transplant within the last 28 days b. Pt is a CMV-seropositive recipient [R+] 2. For: CMV prophylaxis in kidney transplant recipients a. Pt is post kidney transplant within last 7 days b. Pt is high risk (Donor CMV seropositive/Recipient CMV seronegative) 3. Medication will be discontinued on or before 200 days post-transplantation

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

200 days

### **OTHER CRITERIA**

N/A



## **PROLASTIN**

---

### **MEDICATION(S)**

PROLASTIN-C

**PENDING CMS APPROVAL**

## **PROMACTA**

---

### **MEDICATION(S)**

ELTROMBOPAG OLAMINE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For ITP dx: a. Previous failure to corticosteroids, immunoglobulins, OR splenectomy, b. Initial: Evidence of bleeding OR platelet count less than 50,000/microL, c. For Reauth: Platelet count less than 400,000/microL, 2. For Hep C with Thrombocytopenia dx: a. Platelet count less than 75,000/microL, 3. For aplastic anemia dx: a. Pt has an insufficient response to immunosuppressive therapy b. In combination with immunosuppressive therapy for severe aplastic anemia

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

6 Months

### **OTHER CRITERIA**

N/A

## **PURIXAN**

---

### **MEDICATION(S)**

MERCAPTOPURINE 2000 MG/100ML SUSPENSION

**PENDING CMS APPROVAL**

## **PYRUKYND**

---

### **MEDICATION(S)**

PYRUKYND, PYRUKYND TAPER PACK

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Homozygous R479H mutation or 2 non-missense mutations, without the presence of another missense mutation, in the PKLR gene

### **REQUIRED MEDICAL INFORMATION**

1. Initial:

a. Documented pyruvate kinase deficiency (PKD), presence of at least 2 mutant alleles in PKLR gene, of which at least 1 is a missense mutation

b. 6 or more transfusions in the last 12 months

i. If 5 or fewer transfusions, Hb concentration less than or equal to 10.0 g/dL

2. Reauth:

a. Member has increase in Hb compared to baseline and/or reduction in transfusion burden

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

**OTHER CRITERIA**

N/A

## **QINLOCK**

---

### **MEDICATION(S)**

QINLOCK

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of advanced GIST 2. Prior treatment with 3 or more kinase inhibitors, including imatinib

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **QULIPTA**

---

### **MEDICATION(S)**

QULIPTA

**PENDING CMS APPROVAL**

## **RCC**

---

### **MEDICATION(S)**

EVEROLIMUS 10 MG TAB, EVEROLIMUS 2 MG TAB SOL, EVEROLIMUS 2.5 MG TAB, EVEROLIMUS 3 MG TAB SOL, EVEROLIMUS 5 MG TAB, EVEROLIMUS 5 MG TAB SOL, EVEROLIMUS 7.5 MG TAB, TORPENZ

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

For renal angiomyolipoma, requires immediate surgery

### **REQUIRED MEDICAL INFORMATION**

1. For RCC dx: a. Previous failure on either sunitinib or Nexavar, 2. For SEGA or TS dx: a. Patient must require therapeutic intervention and not be a candidate for surgical resection 3. Diagnosis of progressive neuroendocrine tumors of pancreatic origin (PNET) or with well-differentiated, nonfunctional neuroendocrine tumors (NET) of gastrointestinal (GI), or lung origin which are unresectable, locally advanced or metastatic 4. Diagnosed with renal angiomyolipoma with tuberous sclerosis complex with at least one angiomyolipoma greater than or equal to 3cm where there is not an immediate need for surgery 5. Hormone receptor positive HER2-negative breast cancer a. Previous use of one of letrozole or anastrozole b. Use in combination with one of exemastane, tamoxifen, or fulvestrant

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A





## **RETEVMO**

---

### **MEDICATION(S)**

RETEVMO 120 MG TAB, RETEVMO 160 MG TAB, RETEVMO 40 MG TAB, RETEVMO 80 MG TAB

### **PENDING CMS APPROVAL**

## **REVCovi**

---

### **MEDICATION(S)**

REVCovi

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of adenosine deaminase deficiency (ADA) with Severe Combined Immunodeficiency (SCID) phenotype confirmed by one of the following: A. Deficiency of adenosine deaminase (ADA) in plasma, lysed erythrocytes, fibroblasts (cultured from amniotic fluid), or chorionic villus (less than 1% of normal) B. Increase in deoxyadenosine triphosphate (dATP) levels in erythrocyte lysates compared to laboratory standard C. Decrease in ATP concentration in erythrocytes D. Molecular genetic confirmation of mutations in both alleles of the ADA1 gene E. Positive screening by T cell receptor excision circles (TRECs) 2. Not a candidate for or has failed bone marrow transplantation (BMT) 3. Platelets greater than 50,000/cell/microL 4. Reauthorization A. The patient has experienced improvement in their plasma ADA activity, red blood cell dATP levels, immune function, and/or red blood cell dAXP levels

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **REVLIMID**

---

### **MEDICATION(S)**

LENALIDOMIDE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For Multiple Myeloma dx: a. Used in combination with dexamethasone, OR b. As maintenance following autologous hematopoietic stem cell transplantation (auto HSCT) 2. For MCL: a. Pt has previous trial on bortezomib AND pt has trial on at least ONE other previous therapy 3. For transfusion-dependent anemia due to myelodysplastic syndrome a. Low or imitediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q abnormality with or without additional cytogenetic abnormalities 4. For follicular lymphoma and Marginal zone lymphoma (MZL) a. Used in combination with a rituximab product

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **REVUFORJ**

---

### **MEDICATION(S)**

REVUFORJ

**PENDING CMS APPROVAL**

## **REXULTI**

---

### **MEDICATION(S)**

REXULTI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

N/A

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## REYVOW

---

### MEDICATION(S)

REYVOW

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Patient has tried 2 generic triptan medications OR has a contraindication to the use of triptans (e.g. established cardiovascular disease or cerebrovascular disease)

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A

## **REZDIFFRA**

---

### **MEDICATION(S)**

REZDIFFRA

**PENDING CMS APPROVAL**



## REZLIDHIA

---

### MEDICATION(S)

REZLIDHIA

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Diagnosis of relapsed or refractory AML 2. Confirmed IDH-1 mutation

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A

## **REZUROCK**

---

### **MEDICATION(S)**

REZUROCK

**PENDING CMS APPROVAL**

## **ROMVIMZA**

---

### **MEDICATION(S)**

ROMVIMZA

**PENDING CMS APPROVAL**

## **ROZLYTREK**

---

### **MEDICATION(S)**

ROZLYTREK

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of non-small cell lung cancer A. Has reactive oxygen species 1 positive 2. Diagnosis of Neurotrophic receptor tyrosine kinase-positive solid tumors A. Tumor is metastatic or surgical resection likely to result in severe morbidity B. Progression following previous treatment or there is not an adequate alternative treatment 3. For the oral pellets: patient has inability to swallow capsules and solution made from capsules.

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **RUBRACA**

---

### **MEDICATION(S)**

RUBRACA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of recurrent ovarian cancer a. Complete or partial response to platinum-based chemotherapy 2. Rubraca will be used as monotherapy 3. Dx of mCRPC that has deleterious BRCA mutation a. Previous treatment with androgen receptor-directed therapy and a taxane-based chemotherapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **RYDAPT**

---

### **MEDICATION(S)**

RYDAPT

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Used for post-consolidation therapy (maintenance) for dx of AML

### **REQUIRED MEDICAL INFORMATION**

1. For AML dx: a. Cancer is FLT3 mutation positive 2. Dx of systemic mastocytosis a. Systemic mastocytosis is identified as aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **SABRIL**

---

### **MEDICATION(S)**

VIGABATRIN, VIGADRUNE 500 MG PACKET, VIGAFYDE, VIGPODER

### **PENDING CMS APPROVAL**

## **SCEMBLIX**

---

### **MEDICATION(S)**

SCEMBLIX

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Newly diagnosed Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP) 2. Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP): A. Trial and failure of at least 1 TKI 3. Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP) with T315I mutation: A. Documentation of testing for mutation B. Must try and fail ponatinib

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **SIGNIFOR**

---

### **MEDICATION(S)**

SIGNIFOR

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For Cushing's: a. Pt is NOT a candidate for pituitary surgery, b. If pt previously had pituitary surgery: Pt continues to have high 24-hour urinary free cortisol levels c. Patient has tried at least one of the following: i. ketoconazole ii. metyrapone 2. For acromegaly a. Pt has had surgical resection of the pituitary gland OR is not a candidate for surgery/radiation therapy b. Pt has tried at least one of the following: i. bromocriptine ii. cabergoline iii. octreotide acetate c. Pt has tried at least one of the following: octreotide depot or lanreotide

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: Plan year Reauth: Plan Year

### **OTHER CRITERIA**

N/A

## **SIRTURO**

---

### **MEDICATION(S)**

SIRTURO

**PENDING CMS APPROVAL**

## **SOHONOS**

---

### **MEDICATION(S)**

SOHONOS

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of fibrodysplasia ossificans progressiva (FOP) confirmed by genetic testing (documentation must be submitted)

### **AGE RESTRICTION**

Females age 8 years or older and Males age 10 years or older

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **SOMAVERT**

---

### **MEDICATION(S)**

SOMAVERT

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has had surgical resection of the pituitary gland OR is not a candidate for surgery/radiation therapy, 2. Patient has tried at least ONE of the following: a. Bromocriptine, b. Cabergoline, c. Octreotide acetate 3. Patient has tried at least ONE of the following: octreotide depot or lanreotide

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **SPRYCEL**

---

### **MEDICATION(S)**

DASATINIB 100 MG TAB, DASATINIB 140 MG TAB, DASATINIB 20 MG TAB, DASATINIB 50 MG TAB, DASATINIB 70 MG TAB

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase.
2. Treatment for adults with dx of chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy including imatinib.
3. Treatment for adults with dx of Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) with resistance or intolerance to prior therapy.
4. Treatment of pediatric patients 1 year of age and older with dx of Ph+ CML in chronic phase.
5. Treatment of pediatric patients 1 year of age and older with dx of newly diagnosed Ph+ ALL in combination with chemotherapy.

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

**OTHER CRITERIA**

N/A

## **STELARA**

---

### **MEDICATION(S)**

STELARA 45 MG/0.5ML SOLN PRSYR, STELARA 45 MG/0.5ML SOLUTION, STELARA 90 MG/ML SOLN PRSYR, USTEKINUMAB 45 MG/0.5ML SOLN PRSYR, USTEKINUMAB 45 MG/0.5ML SOLUTION, USTEKINUMAB 90 MG/ML SOLN PRSYR

### **PENDING CMS APPROVAL**

## **STIVARGA**

---

### **MEDICATION(S)**

STIVARGA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of metastatic colorectal cancer (CRC)
  - A. Patient has been previously treated with fluoropyrimidine-, oxaliplatin- and irinotecan-based chemotherapy, an anti-VEGF therapy, and, if RAS wild-type, an anti-EGFR therapy.
2. Diagnosis of locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST)
  - A. Patient has been previously treated with imatinib mesylate and sunitinib malate.
3. Diagnosis of hepatocellular carcinoma (HCC)
  - A. Patient has been previously treated with sorafenib

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **SUCRAID**

---

### **MEDICATION(S)**

SUCRAID

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx is confirmed by a Small Bowel Biopsy Disaccharidase Measurement demonstrating 2 SD or more below mean for sucrase activity with or without isomaltase activity

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **SUTENT**

---

### **MEDICATION(S)**

SUNITINIB MALATE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Sutent used as combination therapy with other chemotherapies

### **REQUIRED MEDICAL INFORMATION**

1. Treatment of adult patients with gastrointestinal stromal tumor (GIST)

a. Patient had disease progression on or intolerance to imatinib mesylate.

2. Treatment of adult patients with advanced renal cell carcinoma (RCC).

3. Adjuvant treatment of adult patients at high risk of recurrent RCC following nephrectomy.

4. Treatment of progressive, well-differentiated pancreatic neuroendocrine tumors (pNET) in adult patients with unresectable locally advanced or metastatic disease.

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **SYNAREL**

---

### **MEDICATION(S)**

SYNAREL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

Pt has previous trial on leuprolide acetate

### **AGE RESTRICTION**

1. For CPP: Treatment initiated at or before 8 years of age in girls and 9 years of age in boys 2. For Endometriosis: 18 years old or older

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **SYPRINE**

---

### **MEDICATION(S)**

TRIENTINE HCL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has failure on penicillamine,

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TABLOID**

---

### **MEDICATION(S)**

TABLOID

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Use for chronic lymphocytic leukemia, Hodgkins lymphoma, multiple myeloma, or solid tumors
2. Use in patients whose disease has demonstrated prior resistance to thioguanine

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of   a. Acute Myeloid Leukemia

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TABRECTA**

---

### **MEDICATION(S)**

TABRECTA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of metastatic non-small cell lung cancer (NSCLC) 2. Tumor mutation leading to mesenchymal-epithelial transition (MET) exon 14 skipping

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TADALAFIL**

---

### **MEDICATION(S)**

TADALAFIL 5 MG TAB

**PENDING CMS APPROVAL**

## **TAFINLAR**

---

### **MEDICATION(S)**

TAFINLAR

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1)For unresectable or metastatic melanoma dx: a)Pt is BRAF (V600E or V600K) mutation positive  
2)For adjuvant treatment of melanoma dx: a)Pt is BRAF (V600E or V600K) mutation positive b)Pt has failed Opdivo AND Keytruda 3)For metastatic NSCLC dx: a)Pt is BRAF V600E positive 4)For Locally Advanced or Metastatic Anaplastic Thyroid Cancer dx: a)Pt is BRAF V600E positive with no satisfactory locoregional treatment options 5)For Unresectable or Metastatic Solid Tumors dx: a)Pt is BRAF V600E positive and has progressed following prior treatment and have no satisfactory alternative treatment options 6)For Low-Grade Glioma dx a)Pt is BRAF V600E mutation positive and requires systemic therapy b)Pt has failed Zelboraf (with or without Cotellic) 7)For Tafinlar tablets for oral suspension: Member has inability to swallow capsules

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **TAGRIS**

---

### **MEDICATION(S)**

TAGRIS

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of metastatic non-small cell lung cancer (NSCLC) and will be used as one of the following:  
A. adjuvant therapy after tumor resection, whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations B. locally advanced, unresectable (stage III) NSCLC whose disease has not progressed during or following concurrent or sequential platinum-based chemoradiation therapy and whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations C. first-line treatment of metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations D. first-line treatment of locally advanced or metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations in combination with pemetrexed and platinum-based chemotherapy E. treatment of adult patients with metastatic EGFR T790M mutation positive NSCLC, whose disease has progressed on or after EGFR TKI therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TALTZ**

---

### **MEDICATION(S)**

TALTZ 80 MG/ML SOLN A-INJ, TALTZ 80 MG/ML SOLN PRSYR

### **PA INDICATION INDICATOR**

2 - Some FDA-Approved Indications Only

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Combination therapy with another biologic medication, JAK inhibitor, or Otezla

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Ankylosing Spondylitis (Please note, Taltz is not covered for indications other than Ankylosing spondylitis)
2. Member has tried TWO of: adalimumab (ie Amjevita, Hadlima, etc), Enbrel, or infliximab (ie. Renflexis)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## TALZENNA

---

### MEDICATION(S)

TALZENNA

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Dx of HRR Gene-mutated mCRPC, in combination with enzalutamide (Xtandi) 2. Deleterious or suspected deleterious germline BRCA, HER2-negative locally advanced or metastatic breast cancer

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A

## TARCEVA

---

### MEDICATION(S)

ERLOTINIB HCL

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. For NSCLC dx: a. Pt with EGFR mutation, b. Erlotinib is not used in combination with platinum-based chemotherapy, 2. For pancreatic cancer dx: a. Combination with gemcitabine

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A

## **TARGRETIN**

---

### **MEDICATION(S)**

BEXAROTENE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. If female: Pt planning to become pregnant

### **REQUIRED MEDICAL INFORMATION**

1. Capsules: a. Pt has previous failure on at least ONE of the following: i. Antineoplastic chemotherapy, ii. Interferon alfa and gamma, iii. Interleuking-12, iv. Interleukin-2, 2. Gel: a. Pt has previous failure on at least ONE of the following: i. PUVA, ii. UVB, iii. EVT, iv. Photophoresis, v. Systemic cytotoxic chemotherapy, vi. Topical nitrogen mustard, vii. Topical carmustine

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TARPEYO**

---

### **MEDICATION(S)**

FILSPARI, TARPEYO

### **PENDING CMS APPROVAL**

## **TASIGNA**

---

### **MEDICATION(S)**

NILOTINIB HCL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

For Ph+ CML Dx: Pts who have BCR-ABL1 mutations T315I, Y253H, E255K/V, F359V/C/I, or G250E

### **REQUIRED MEDICAL INFORMATION**

1. Adult or pediatric with newly diagnosed Philadelphia chromosome positive (Ph+) CML in chronic phase 2. Adult with chronic phase and accelerated phase Ph+ CML a. Resistant or intolerant to imatinib 3. Pediatric patient with chronic phase or accelerated phase Ph+ CML a. Resistant or intolerant to prior TKI therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

# **TAVALISSE**

---

## **MEDICATION(S)**

TAVALISSE

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

N/A

## **REQUIRED MEDICAL INFORMATION**

1. Initial: a. Pt has previous trial on previous therapy of at least ONE of the following: i. Corticosteroids ii. Immunoglobulins iii. Splenectomy iv. Thrombopoietin Receptor Agonist b. Platelet count is less than 50000/micoL 2. Reauth: a. Platelet count is greater than 50000/micoL

## **AGE RESTRICTION**

N/A

## **PRESCRIBER RESTRICTION**

N/A

## **COVERAGE DURATION**

Initial: 3 Months Reauth: Plan Year

## **OTHER CRITERIA**

N/A



## **TAVNEOS**

---

### **MEDICATION(S)**

TAVNEOS

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Patient currently requires dialysis

### **REQUIRED MEDICAL INFORMATION**

Initial: 1. Diagnosis of ANCA-associated vasculitis 2. ANCA-antibody titer test 3. BVAS score (1 major item, 3 non-major items, or 2 renal items of proteinuria and hematuria) 4. Patient is currently receiving rituximab or cyclophosphamide Reauth: 1. Patient has experienced improvement in BVAS score

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Initial: 6 months Reauth: Plan Year

### **OTHER CRITERIA**

N/A

## **TAZVERIK**

---

### **MEDICATION(S)**

TAZVERIK

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx: Metastatic or locally advanced epithelioid sarcoma a. Diagnosis confirmed by pathology b. SMARCB1/INI1 deficient tumor c. Tumor is not eligible for complete resection 2. Dx: Follicular lymphoma a. Relapsed or refractory tumor i. positive for EZH2 mutation ii. Previous use of at least 2 prior systemic therapies b. No satisfactory alternative treatment options available

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TEPMETKO**

---

### **MEDICATION(S)**

TEPMETKO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of metastatic non-small cell lung cancer (NSCLC) harboring mesenchymal-epithelial transition (MET) exon 14 skipping alterations.

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

# **TIBSOVO**

---

## **MEDICATION(S)**

TIBSOVO

## **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

## **OFF LABEL USES**

N/A

## **EXCLUSION CRITERIA**

N/A

## **REQUIRED MEDICAL INFORMATION**

1. Treatment of newly diagnosed AML in adults 75 years or older, or who have comorbidities that preclude the use of intensive induction chemotherapy and will be used either in combination with azacitadine or as monotherapy
2. Diagnosis of relapsed or refractory AML.
3. Diagnosis of relapsed or refractory myelodysplastic syndromes.
4. Diagnosis of locally advanced or metastatic cholangiocarcinoma who have been previously treated

## **AGE RESTRICTION**

N/A

## **PRESCRIBER RESTRICTION**

N/A

## **COVERAGE DURATION**

Plan Year

## **OTHER CRITERIA**

N/A

## **TRIKAFTA**

---

### **MEDICATION(S)**

TRIKAFTA

**PENDING CMS APPROVAL**

## TRUQAP

---

### MEDICATION(S)

TRUQAP

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

1. diabetes mellitus type 1, type 2, requiring insulin treatment or HbA1c greater than or equal to 8%

### REQUIRED MEDICAL INFORMATION

1. Dx of HR-positive, HER2-negative, locally advanced or metastatic breast cancer 2. Patient has PIK3CA/AKT1/PTEN-alterations. 3. Patient has failed at least one endocrine-based regimen in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy 4. Truqap with be used in combination with fulvestrant

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan Year

### OTHER CRITERIA

N/A

# TUKYSA

---

## MEDICATION(S)

TUKYSA

## PA INDICATION INDICATOR

1 - All FDA-Approved Indications

## OFF LABEL USES

N/A

## EXCLUSION CRITERIA

N/A

## REQUIRED MEDICAL INFORMATION

1. Diagnosis of advanced unresectable or metastatic HER2-positive breast cancer, including patients with brain metastases:

- a. Pt has received one or more prior anti-HER2-based regimens in the metastatic setting
- b. Tukysa will be used in combination with trastuzumab and capecitabine.

2. Diagnosis of RAS wild-type HER2-positive unresectable or metastatic colorectal cancer

- a. Cancer has progressed following treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy,
- b. Tukysa will be used in combination with trastuzumab

## AGE RESTRICTION

N/A

## PRESCRIBER RESTRICTION

N/A

## COVERAGE DURATION

Plan Year

**OTHER CRITERIA**

N/A



## **TURALIO**

---

### **MEDICATION(S)**

TURALIO 125 MG CAP

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

A. Diagnosis of Tenosynovial Giant Cell Tumor B. Condition is associated with severe morbidity or functional limitations C. Surgery will NOT improve status

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TYENNE**

---

### **MEDICATION(S)**

TYENNE 162 MG/0.9ML SOLN A-INJ, TYENNE 162 MG/0.9ML SOLN PRSYR

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

1. Combination use with another biologic medications, janus kinase inhibitor (JAK), or Otezla

### **REQUIRED MEDICAL INFORMATION**

1. Dx of: A. Cytokine release syndrome (CRS) B. Giant cell arteritis (GCA) C. Systemic juvenile idiopathic arthritis (JIA) D. Moderate to severe rheumatoid arthritis (RA) E. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) F. Covid-19 2, For RA or JIA A. Pt has failed at least three months therapy on at least ONE of the following: i. methotrexate, ii. leflunomide, iii. hydroxychloroquine, iv. sulfasalazine, OR contraindication to use (Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease, Pregnancy, breastfeeding, or currently planning pregnancy, Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia), Elevated liver transaminases, Hypersensitivity, or history of intolerance or adverse event, Interstitial pneumonitis or clinically significant pulmonary fibrosis, Myelodysplasia, Renal impairment

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **TYKERB**

---

### **MEDICATION(S)**

LAPATINIB DITOSYLATE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For advanced or metastatic HER-2 positive breast cancer dx: a. Previous failure on anthracycline, taxane, and trastuzumab AND b. Combination therapy with capecitabine 2. For postmenopausal HER-2 receptor hormone receptor positive breast cancer dx: a. Combination therapy with aromatase inhibitor

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **UBRELVY**

---

### **MEDICATION(S)**

UBRELVY

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Patient has tried 2 generic triptan medications OR has a contraindication to the use of triptans (e.g. established cardiovascular disease or cerebrovascular disease)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **UPTRAVI**

---

### **MEDICATION(S)**

UPTRAVI 1000 MCG TAB, UPTRAVI 1200 MCG TAB, UPTRAVI 1400 MCG TAB, UPTRAVI 1600 MCG TAB, UPTRAVI 200 & 800 MCG TAB THPK, UPTRAVI 200 MCG TAB, UPTRAVI 400 MCG TAB, UPTRAVI 600 MCG TAB, UPTRAVI 800 MCG TAB

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Initial:

a. If pt has a positive vasoreactivity test:

i. Pt has failed maximum tolerated doses of calcium channel blockers,

b. Pt has previous trial on at least ONE of the following:

i. ambrisentan, ii. bosentan iii. Opsumit iv. Adempas

2. Reauth:

a. Pt has been reassessed within the past 6 months.

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Pulmonologist or Cardiologist

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **USTEKINUMAB**

---

### **MEDICATION(S)**

PYZCHIVA 45 MG/0.5ML SOLN PRSYR, PYZCHIVA 45 MG/0.5ML SOLUTION, PYZCHIVA 90 MG/ML SOLN PRSYR, SELARSDI 45 MG/0.5ML SOLN PRSYR, SELARSDI 90 MG/ML SOLN PRSYR

### **PENDING CMS APPROVAL**

## VALCHLOR

---

### MEDICATION(S)

VALCHLOR

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Pt has previous trial on at least ONE previous skin directed therapy of the following: a. Topical corticosteroid, b. Topical carmustine, c. Topical retinoid, d. Radiation therapy, e. Phototherapy

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A

## **VANFLYTA**

---

### **MEDICATION(S)**

VANFLYTA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Newly diagnosed acute myeloid leukemia that is FLT3 internal tandem duplication (ITD)-positive as detected by an FDA approved test. 2. Will be taken in combination with standard cytarabine and anthracycline induction, cytarabine consolidation, and as maintenance monotherapy following consolidation chemotherapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



## **VAXCHORA**

---

### **MEDICATION(S)**

VAXCHORA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Member is traveling to area of active cholera transmission (Please note, the dates and location of active cholera transmission area member is going to need to be documented in request) 2. Member is between the ages of 2 and 64

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **VELTASSA**

---

### **MEDICATION(S)**

VELTASSA

**PENDING CMS APPROVAL**

## **VEMLIDY**

---

### **MEDICATION(S)**

VEMLIDY

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

Pt has decompensated hepatic impairment

### **REQUIRED MEDICAL INFORMATION**

Pt has previous trial on entecavir

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **VENCLEXTA**

---

### **MEDICATION(S)**

VENCLEXTA, VENCLEXTA STARTING PACK

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For CLL/SLL dx 2. For AML dx: a. Pt is ineligible for induction therapy OR b. Pt is 75 years or older

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **VERQUVO**

---

### **MEDICATION(S)**

VERQUVO

**PENDING CMS APPROVAL**

## **VERZENIO**

---

### **MEDICATION(S)**

VERZENIO

**PENDING CMS APPROVAL**

## **VIJOICE**

---

### **MEDICATION(S)**

VIJOICE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Confirmed/documented diagnosis of PIK3CA Related Overgrowth Spectrum (PROS) a. Patient has mutation in the PIK3CA gene.

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **VIMKUNYA**

---

### **MEDICATION(S)**

VIMKUNYA

**PENDING CMS APPROVAL**



## VITRAKVI

---

### MEDICATION(S)

VITRAKVI

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. solid tumor with a NTRK gene fusion 2. Metastatic or unable to have surgery 3. Received previous treatment

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A

## **VIVJOA**

---

### **MEDICATION(S)**

VIVJOA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Hx of recurrent vulvovaginal candidiasis, defined as at least 3 acute episodes in the last 12 months  
2. Patient must be one of the following: A. Post-menopausal B. Not of reproductive potential (i.e. tubal ligation, hysterectomy, etc)  
3. Patient has experienced a recurrence during or following 6 months of oral fluconazole maintenance therapy

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

14 weeks

### **OTHER CRITERIA**

N/A

## **VIVOTIF**

---

### **MEDICATION(S)**

VIVOTIF

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Member meets one of the following:

A. Traveling to an area with recognized risk of exposure to *S. typhi* (dates and location of the area member is going to need to be documented in request)

B. Has intimate exposure to a *S. typhi* carrier (documentation must be provided with request)

C. a microbiology laboratorian frequently working with *S. typhi* (documentation must be provided with request)

2. Member has not received vaccination for *S. typhi* in the last 5 years

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

1 Month

### **OTHER CRITERIA**

N/A



## **VIZIMPRO**

---

### **MEDICATION(S)**

VIZIMPRO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1) First line therapy 2) EGFR exon 19 deletion or EFGR exon 21 L858R substitution

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## VONJO

---

### MEDICATION(S)

VONJO

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Diagnosis of intermediate or high-risk primary or secondary myelofibrosis (MF) a. For secondary MF: post-polycythemia vera or post-essential thrombocythemia 2. Documentation showing platelet counts below 50,000/mm<sup>3</sup> within the last 30 days 3. For intermediate risk: Inadequate response or intolerance to hydroxyurea, Pegasys, or Jakafi 4. For high risk: patient is not a candidate for transplant 5. Reauthorization: CBC and platelet count required. If above 50,000mm<sup>3</sup>, Vonjo is no longer indicated. Jakafi is approved for use in patients with platelet counts above 50,000/mm<sup>3</sup>

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Initial: 6 months Reauthorization: Plan Year

### OTHER CRITERIA

N/A

## **VORANIGO**

---

### **MEDICATION(S)**

VORANIGO

**PENDING CMS APPROVAL**

## **VOSEVI**

---

### **MEDICATION(S)**

VOSEVI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Confirmation of genotype

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

12 Weeks

### **OTHER CRITERIA**

N/A



## **VOTRIENT**

---

### **MEDICATION(S)**

PAZOPANIB HCL

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Soft Tissue Sarcoma dx: a. Previous trial on at least ONE prior therapy 2. Advanced renal cell carcinoma dx

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **VOXZOGO**

---

### **MEDICATION(S)**

VOXZOGO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of achondroplasia, confirmed by genetic testing 2. Ambulatory and able to stand without assistance 3. Member has open epiphyses 4. Reauthorization a) No evidence of growth plate closure (proximal tibia, distal femur)

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **VYNDAQEL**

---

### **MEDICATION(S)**

VYNDAMAX, VYNDAQEL

### **PENDING CMS APPROVAL**

## **WELIREG**

---

### **MEDICATION(S)**

WELIREG

**PENDING CMS APPROVAL**

## **WINREVAIR**

---

### **MEDICATION(S)**

WINREVAIR

**PENDING CMS APPROVAL**

## **XALKORI**

---

### **MEDICATION(S)**

XALKORI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1) Dx with: a) metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK) b) metastatic non-small cell lung cancer (NSCLC) whose tumors are ROS1-positive c) relapsed or refractory, systemic anaplastic large cell lymphoma (ALCL) that is ALK-positive d) unresectable, recurrent, or refractory ALK-positive inflammatory myofibroblastic tumor 2) For NSCLC and ALK-positive inflammatory myofibroblastic tumor: Patient has failed/intolerance/contraindication to Alecensa

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan year

### **OTHER CRITERIA**

N/A

## **XDEMVY**

---

### **MEDICATION(S)**

XDEMVY

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of Demodex blepharitis

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

Ophthalmologist

### **COVERAGE DURATION**

6 weeks

### **OTHER CRITERIA**

N/A

## **XELJANZ**

---

### **MEDICATION(S)**

XELJANZ, XELJANZ XR

### **PENDING CMS APPROVAL**



## **XENAZINE**

---

### **MEDICATION(S)**

TETRABENAZINE

### **PA INDICATION INDICATOR**

4 - All FDA-Approved Indications, Some Medically-Accepted Indications

### **OFF LABEL USES**

Tardive Dyskinesia

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of chorea due to Huntington's Disease OR 2. Diagnosis of Tardive Dyskinesia

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **XERMELO**

---

### **MEDICATION(S)**

XERMELO

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Initial:

- a. Pt has more than 4 bowel movements a day despite treatment with sandostatin analog therapy for at least 3 months
- b. Pt has previous trial of lomotil AND loperamide
- c. Xermelo will be used in combination with ocreotide depot or lanreotide

2. Reauth:

- a. Pt has experienced improvement in bowel movement frequency since starting Xermelo

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

6 Months

### **OTHER CRITERIA**

N/A

## **XHANCE**

---

### **MEDICATION(S)**

XHANCE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1.Diagnosis of Chronic Rhinosinusitis with or without Nasal Polyps. 2. Previous use of mometasone intranasal spray

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **XIFAXAN**

---

### **MEDICATION(S)**

XIFAXAN

### **PA INDICATION INDICATOR**

4 - All FDA-Approved Indications, Some Medically-Accepted Indications

### **OFF LABEL USES**

1. Clostridium difficile infection

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of: a. Clostridium difficile (C. diff.) infection b. Hepatic encephalopathy (HE) c. Irritable bowel syndrome (IBS) d. Traveler's diarrhea 2. Diagnosis of Hepatic encephalopathy (HE) a. Previous failure on or has intolerance to lactulose therapy 3. Diagnosis of Irritable bowel syndrome (IBS) a. Previous failure of at least TWO antispasmodic or antibiotic treatments (e.g., amoxicillin-clavulanate, cephalexin, ciprofloxacin, dicyclomine, doxycycline, gentamicin, metronidazole, neomycin, trimethoprim-sulfamethoxazole) 4. Diagnosis of Clostridium difficile (C. diff) infection a. Patient has experienced relapse after prior use of oral vancomycin

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **XOLAIR**

---

### **MEDICATION(S)**

XOLAIR

**PENDING CMS APPROVAL**

## **XOSPATA**

---

### **MEDICATION(S)**

XOSPATA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Relapsed or refractory AML 2. patient has a FLT3 mutation detected by an FDA-approved test

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **XPOVIO**

---

### **MEDICATION(S)**

XPOVIO (100 MG ONCE WEEKLY) 50 MG TAB THPK, XPOVIO (40 MG ONCE WEEKLY) 10 MG TAB THPK, XPOVIO (40 MG ONCE WEEKLY) 40 MG TAB THPK, XPOVIO (40 MG TWICE WEEKLY) 40 MG TAB THPK, XPOVIO (60 MG ONCE WEEKLY) 60 MG TAB THPK, XPOVIO (60 MG TWICE WEEKLY), XPOVIO (80 MG ONCE WEEKLY) 40 MG TAB THPK, XPOVIO (80 MG TWICE WEEKLY)

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Diagnosis of multiple myeloma:

A. Patient has received at least one prior therapy

B. Xpovio will be used in combination with bortezomib and dexamethasone

2. Diagnosis of relapsed or refractory multiple myeloma:

A. Patient has received at least four prior therapies

B. Patient's disease is refractory to at least two proteasome inhibitors, at least two immunomodulatory agents, and an anti-CD38 monoclonal antibody

C. Xpovio will be used in combination with dexamethasone.

3. Diagnosis of relapsed or refractory diffuse large B-cell lymphoma (DLBCL), not otherwise specified, including DLBCL arising from follicular lymphoma

A. Patient has previous use of at least 2 lines of systemic therapy.



**AGE RESTRICTION**

N/A

**PRESCRIBER RESTRICTION**

N/A

**COVERAGE DURATION**

Plan Year

**OTHER CRITERIA**

N/A

## **XTANDI**

---

### **MEDICATION(S)**

XTANDI

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1) Diagnosis of non-metastatic castration sensitive prostate cancer (nmCSPC) a)Patient has failed androgen deprivation therapy (ADT) with one of the following: orchiectomy, LHRH agonist (leuprolide, Lupron, Zoladex), LHRH antagonist (Firmagon) 2)Diagnosis of metastatic castration-sensitive prostate cancer (mCSPC) a)Patient has failed ADT and abiraterone 250mg 3)Diagnosis of non-metastatic castration-resistant prostate cancer (nmCRPC) a)Patient has PSA doubling time of less than 10 months

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **YORVIPATH**

---

### **MEDICATION(S)**

YORVIPATH

**PENDING CMS APPROVAL**

## **ZARXIO**

---

### **MEDICATION(S)**

NEUPOGEN, RELEUKO 300 MCG/0.5ML SOLN PRSYR, RELEUKO 480 MCG/0.8ML SOLN PRSYR, ZARXIO

**PENDING CMS APPROVAL**

## **ZEJULA**

---

### **MEDICATION(S)**

ZEJULA 100 MG TAB, ZEJULA 200 MG TAB, ZEJULA 300 MG TAB

### **PENDING CMS APPROVAL**

## **ZELBORAF**

---

### **MEDICATION(S)**

COTELLIC, ZELBORAF

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. For Metastatic Melanoma dx: a. Pt is BRAF V600E positive for Zelboraf monotherapy OR b. BRAF V600E or V600K positive for Zelboraf plus Cotellic 2. For Erdheim-Chester Disease: a. Zelboraf monotherapy b. Pt is BRAF V600 positive

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ZOLINZA**

---

### **MEDICATION(S)**

ZOLINZA

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Pt has progressive, persistent, or recurrent disease, 2. Pt has tried at least TWO prior systemic therapies

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## **ZONISADE**

---

### **MEDICATION(S)**

ZONISADE

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. Dx of partial-onset seizures 2. Inability to swallow tablets and capsules

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A



**ZTALMY**

---

**MEDICATION(S)**

ZTALMY

**PENDING CMS APPROVAL**

## **ZURZUVAE**

---

### **MEDICATION(S)**

ZURZUVAE

**PENDING CMS APPROVAL**

## **ZYDELIG**

---

### **MEDICATION(S)**

ZYDELIG

### **PA INDICATION INDICATOR**

1 - All FDA-Approved Indications

### **OFF LABEL USES**

N/A

### **EXCLUSION CRITERIA**

N/A

### **REQUIRED MEDICAL INFORMATION**

1. CLL dx: a. Used in combination with rituximab

### **AGE RESTRICTION**

N/A

### **PRESCRIBER RESTRICTION**

N/A

### **COVERAGE DURATION**

Plan Year

### **OTHER CRITERIA**

N/A

## ZYKADIA

---

### MEDICATION(S)

ZYKADIA

### PA INDICATION INDICATOR

1 - All FDA-Approved Indications

### OFF LABEL USES

N/A

### EXCLUSION CRITERIA

N/A

### REQUIRED MEDICAL INFORMATION

1. Dx with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK)-positive 2. Patient has failed/intolerance/contraindication to Alecensa

### AGE RESTRICTION

N/A

### PRESCRIBER RESTRICTION

N/A

### COVERAGE DURATION

Plan year

### OTHER CRITERIA

N/A